



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification<sup>6</sup>:C12N 15/31, C07K 14/395, G01N 33/68,  
C12N 15/62, C12Q 1/00, C12N 1/16 //  
(C12N 1/16, C12R 1:865)

A1

(11) International Publication Number:

WO 96/12806

(43) International Publication Date:

2 May 1996 (02.05.96)

(21) International Application Number: PCT/US95/13580

(22) International Filing Date: 23 October 1995 (23.10.95)

(30) Priority Data:

08/328,322

24 October 1994 (24.10.94)

US

(71) Applicant: THE BOARD OF TRUSTEES OF THE LELAND  
STANFORD JUNIOR UNIVERSITY [US/US]; Stanford,  
CA 94305 (US).(72) Inventors: HUANG, Laiqiang; 365 Chiquita Avenue, #8,  
Mountain View, CA 94041 (US). CYERT, Martha, S.; 590  
Ashton Avenue, Palo Alto, CA 94306 (US).(74) Agent: SHOLTZ, Charles, K.; Dehlinger & Associates, P.O.  
Box 60850, Palo Alto, CA 94306-1546 (US).(81) Designated States: AU, CA, JP, European patent (AT, BE,  
CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT,  
SE).

## Published


With international search report.

Before the expiration of the time limit for amending the  
claims and to be republished in the event of the receipt of  
amendments.

(54) Title: CALCINEURIN INTERACTING PROTEIN COMPOSITIONS AND METHODS

## (57) Abstract

An identification and characterization of a calcineurin interacting (CNI) protein effective to enhance immunosuppressive effects of calcineurin-targeted immunosuppressants by potentiating an interaction of an immunophilin with calcineurin is described herein. One embodiment of the invention is the CNI polypeptide encoded by the *CNI* gene of *Saccharomyces cerevisiae*. Polynucleotides encoding a CNI protein are also described. Also described are yeast cells carrying mutations in the *CNI* gene. Further, a method of identifying a small molecule immunosuppressant compound is described. The methods include the use of a cell-based two hybrid protein-protein interaction assay, wherein one of two fusion hybrid proteins in a cell contains a subunit of calcineurin, and the other of two fusion hybrid proteins contains an immunophilin.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**CALCINEURIN INTERACTING PROTEIN**  
**COMPOSITIONS AND METHODS**

**Field of the Invention**

- 5       The present invention relates to compounds affecting the function of calcineurin, particularly interactions of calcineurin with immunosuppressant drugs.

**References**

- Ausubel, F.M., *et al.*, *Current Protocols in Molecular Biology* (John Wiley and Sons, Inc., Media, PA).
- 10       Bartel, P., *et al.*, *BioTechniques* 14:920-924 (1993).
- Beach, D., *et al.*, *Mol. Gen. Genet.* 187:326-329 (1982).
- Better, M., *et al.*, *Science* 240:1041-1043 (1988).
- Breeden, J., *et al.*, *Cold Spring Harbor Symp. Quant. Biol.* 50:643 (1985).
- 15       Brent, R., *et al.*, *Cell*, 43:729-736 (1985).
- Briggs J.D., *Immunology Letters*, 29(1-2):89-94 (1991).
- Cabilly, S., *et al.*, European Patent Application No. 84/302368.0, Publication No. 0 125 023, published 14 November 1984.
- Chien, C.-t., *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 88:9578 (1991).
- 20       Clipstone, N.A., *et al.*, *Nature* 357:695 (1992).
- Cyert, M.S., *Current Biology* 2:18 (1992).
- Cyert, M.S., *et al.*, *PNAS* 88:7376 (1991).
- Cyert, M.S., *et al.*, *Mol. Cell. Biol.* 12:3460 (1992).
- Dayhoff, M.O., in Atlas of Protein Sequence and Structure, National Biomedical
- 25       Research Foundation 5:101-110, and Supplement 2 to this volume, pp. 1-10.
- Dice, J.F., *FASEB J.* 1:349-357 (1987).
- Durfee, T., *et al.*, *Genes & Development* 7:555 (1993).
- Fields, S., *et al.*, *Nature* 340:245 (1989).
- Foor, F., *et al.*, *Nature* 360:682 (1992).
- 30       Guerini, D., *et al.*, *Adv. Prot. Phosphatases* 6:391 (1991).
- Guthrie, C., *et al.*, *Meth. Enzymol.* 194:1-774 (1992).
- Gyuris J., *et al.*, *Cell*, 75:791-803 (1993).
- Harlow, E., *et al.*, Antibodies: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, NY (1988).
- 35       Heitman, J., *et al.*, *PNAS* 88:1948-52 (1991).
- Hill, J.E., *et al.*, *Yeast* 2:163-67 (1986).

- Kennedy, M.S., et al., *Am. J. Med.* 78:978 (1983).
- Klee, C.B., et al., "The calmodulin-regulated protein phosphatase" In *Calmodulin*, Klee, C.P., ed. (Elsevier, Amsterdam) (1988).
- Kunkle, T.A., *PNAS* 82:488-92 (1985).
- 5 Laemmli, U., *Nature* 227:680-85 (1970).
- Liu, J., et al., *Cell* 66:807 (1991).
- Ma, J., et al., *Cell* 55:443-446 (1988).
- Platt, J.L., et al., *Immunology Today*, 11(12):450 (1990).
- Reneke, J.E., et al., *Cell* 55:221 (1988).
- 10 Roberts, J.P., et al., *Ann. Rev. Med.*, 40:287 (1989).
- Rogers, et al., *Science* 234:364 (1986).
- Rose, M.D., et al., *Methods in Yeast Genetics* (Cold Spring Harbor, NY, 1990).
- Sambrook, J., et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor, NY, 1972).
- 15 Sanger, F., et al., *PNAS* 74:5463-67 (1977).
- Schiestl, R.H., et al., *Curr. Genet.* 16:339 (1989).
- Schreiber, S.L., *Science* 251:283 (1991).
- Schreiber, S.L., et al., *Immunol. Today* 13:136 (1992).
- Sherman, F., et al., *Methods in Yeast Genetics* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY) (1979).
- 20 Sikorski, R.S., et al., *Genetics* 122:19-27 (1989).
- Skerra, A., et al., *Science* 240:1038-1041 (1988).
- Storb, R., *Blood* 66:698 (1985).
- Storb, R., et al., *N. Engl. J. Med.* 314:729 (1986).
- 25 Triezenberg, S.J., et al., *Genes & Development*, 2:718-29 (1988).
- Weiss, A., et al., *Cell* 76:263 (1994).
- Wilson, I., et al., *Cell* 37:767 (1984).
- Yang, X., et al., *Science* 257:680 (1992).

30 **Background of the Invention**

The immune system functions as the body's major defense against diseases caused by invading organisms. This complex system fights disease by killing invaders such as bacteria, viruses, parasites or cancerous cells while leaving the body's normal tissues unharmed. The immune system's ability to distinguish the body's normal tissues, or self, from foreign or

cancerous tissue, or non-self, is an essential feature of normal immune system function. A second essential feature is memory, the ability to remember a particular foreign invader and to mount an enhanced defensive response when the previously encountered invader returns. The loss of recognition of a particular tissue as self and the subsequent immune response directed against that tissue produce serious illness.

An autoimmune disease results from the immune system attacking the body's own organs or tissues, producing a clinical condition associated with the destruction of that tissue. An autoimmune attack directed against the joint lining tissue results in rheumatoid arthritis; an attack against the conducting fibers of the nervous system results in multiple sclerosis. The autoimmune diseases most likely share a common pathogenesis and the need for safe and effective therapy. One type of therapy that has been employed in combating autoimmune disease is treatment with immunosuppressant drugs, such as cyclosporin A, FK506 and rapamycin. While the treatments are often effective, the drugs typically have undesirable side effects, including neurotoxicity, nephrotoxicity, hypertension, and metabolic disorder. Many of these side effects are due to the drugs' action on cells other than those of the immune system.

In addition to their use in treating autoimmune conditions, immunosuppressive agents have also been used in treating or preventing transplantation rejection. Organ transplantation involving human organ donors and human recipients (allogeneic grafts), and non-human primate donors and human recipients (xenogeneic grafts), has received considerable medical and scientific attention (*e.g.*, Roberts, 1989; Platt, 1990). To a great extent, this effort has been aimed at eliminating, or at least reducing, the problem of rejection of the transplanted organ. In the absence of adequate immunosuppressive therapy, the transplanted organ is destroyed by the host immune system.

Presently, the most commonly used agents for preventing transplant rejection include corticosteroids, cytotoxic drugs that specifically inhibit T cell activation such as azathioprine, immunosuppressive drugs such as cyclosporin A, and specific antibodies directed against T lymphocytes or surface receptors that mediate their activation (Briggs, 1991; Kennedy, 1983; Storb, 1985; Storb, 1986). All of these drug therapies are limited in effectiveness, in part because the doses needed for effective treatment of transplant rejection may increase the patient's susceptibility to infection by a variety of opportunistic invaders, and in part because of direct toxicity and other side effects.

Cyclosporin A, currently the most effective and most commonly used agent, is significantly toxic to the kidney. This nephrotoxicity limits the quantity of drug that can be

safely given. The physician is frequently forced to administer sub-optimal doses of the drug because of this toxicity. A preparation capable of potentiating the action of immunosuppressive agents such as cyclosporin A on the immune system, thus allowing the administration of lower doses of drug, would be of considerable value in reducing the morbidity and mortality associated with transplantation.

### **Summary of the Invention**

In one embodiment, the present invention includes polypeptide compositions effective to enhance immunosuppressive effects of calcineurin-targeted immunosuppressants by potentiating an interaction of an immunophilin with calcineurin. The present invention includes the isolation and characterization of a calcineurin interacting protein, CNI, having these properties. Also disclosed herein are methods for the isolation and characterization of further CNI-related sequences and sequences of CNI-variants. The amino acid sequences presented as SEQ ID NO:2 and SEQ ID NO:5 are exemplary of the polypeptides of the present invention.

The present invention also includes a CNI polypeptide fragment that interacts specifically with the "A" subunits of calcineurin (CNA1 and CNA2), but not with calcineurin "B" subunit (CNB1). In one embodiment, this fragment has an amino acid sequence of between 15 and 915 amino acids in length, for example, the c-terminal 306 amino acids of the CNI protein (CNIc).

Included aspects of the invention are an CNI polypeptide; a recombinant CNI polypeptide; and a fusion polypeptide comprised of an CNI polypeptide. Exemplary fusion proteins include fusions to  $\beta$ -galactosidase.

The invention further includes isolated nucleic acid sequences encoding the above described polypeptides and polypeptide fragments. Exemplary nucleic acid sequences include the sequences presented as SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4 and SEQ ID NO:6. The present invention includes CNI-encoding genomic polynucleotides, cDNAs thereto and complements thereof. With respect to polynucleotides, some aspects of the invention include: a purified CNI-encoding genomic polynucleotide; CNI polypeptide-encoding RNA and DNA polynucleotides; recombinant CNI polypeptide-encoding polynucleotides; a recombinant vector including any of the above recombinant polynucleotides, and a host cell transformed with any of these vectors. Another aspect of the invention is a polynucleotide probe for CNI polypeptide-encoding sequences.

Portions of a CNI-polypeptide coding sequences are effective as probes to isolate variants coding sequences which occur naturally, or to determine the presence of such coding

sequences in nucleic acid samples. Such probes include hybridization screening probes and polymerase chain reaction amplification primers specific for CNI-polypeptide coding sequences. Homologues of CNI may be isolated from a number of sources, such as other types of yeast cells (*e.g.*, *Schizosaccharomyces*) or mammalian cells (*e.g.*, human).

5 Other aspects of the invention include: a recombinant expression system which incorporates an open reading frame (ORF) derived from CNI polypeptide-encoding sequences, wherein the ORF is linked operably to a control sequence which is compatible with a desired host, a cell transformed with the recombinant expression system, and a polypeptide produced by the transformed cell. Typically the expression system includes a vector having (a) a nucleic  
10 acid containing an open reading frame that encodes a CNI-polypeptide; and (b) regulatory sequences effective to express said open reading frame in a host cell. The regulatory sequence may include sequences useful for targeting or secretion of the CNI-polypeptide: such as a secretory signal recognized in yeast or bacterial expression systems.

The invention includes a method of recombinantly producing CNI-polypeptides. In the  
15 method, a recombinant expression system containing an open reading frame (ORF) having a polynucleotide sequence which encodes a CNI-polypeptide, where the vector is designed to express the ORF in the host, is introduced into suitable host cells. The host is then cultured under conditions resulting in the expression of the ORF sequence. The CNI-polypeptide sequences discussed above are examples of suitable CNI-polypeptides. Numerous vectors and  
20 their corresponding hosts are useful in the practice of this method of the invention, including, but not limited, to the vectors described herein for expression in yeast cells, and lambda gt11 phage vector and *E. coli* cells. Other host cells include insect and mammalian cell expression systems.

The invention also includes purified antibodies that are immunoreactive with a CNI-  
25 polypeptide. The antibodies may be polyclonal or monoclonal. Antibodies that are specifically immunoreactive with CNI-polypeptides may be useful for the isolation of CNI-polypeptide homologues from other cell type sources (*e.g.*, mammalian).

The present invention also includes, a method of identifying a small molecule immunosuppressant compound. In the method, a cell-based two hybrid protein-protein  
30 interaction assay is constructed where one of two fusion hybrid proteins in the cell contains a subunit of calcineurin, and the other of two fusion hybrid proteins contains an immunophilin. The cell is then contacted with the small molecule being tested. A small molecule is identified as an immunosuppressant if the molecule potentiates an interaction between the two hybrid proteins. In one embodiment, the method is carried out using yeast cells, where one of the two

fusion hybrid proteins contains a GAL4 activation domain and the other of two fusion hybrid proteins contains a GAL4 binding domain. The method can be carried out where subunits of calcineurin are from any cell source, in particular, yeast or mammalian cells (including human cells). The subunit may, for example, be yeast calcineurin subunit CNA1 or CNA2, or human calcineurin subunit "A". The immunophilin can, for example, be cyclophilins or FK506-binding proteins (*e.g.*, FKBP12) typically from a homologous cell source.

Also included in the present invention is another method of identifying a small molecule immunosuppressant compound. In the method, a cell-based two hybrid protein-protein interaction assay is constructed, wherein one of two fusion hybrid proteins in a cell contains an "A" subunit of calcineurin, and the other of two fusion hybrid proteins contains a CNI polypeptide. The cell preferably, but not necessarily, also contains a vector construct causing overexpression, or increased expression, of a "B" subunit of calcineurin. The cell is then contacted with the small molecule being tested. A small molecule is identified as an immunosuppressant if the molecule potentiates an interaction between the two hybrid proteins.

This method is used to identify compounds (like FK506) that potentiate the interaction between CNI and CNA1. In one embodiment, the method is carried out using yeast cells, where one of the two fusion hybrid proteins contains a GAL4 activation domain and the other of two fusion hybrid proteins contains a GAL4 binding domain. The method can be carried out where subunits of calcineurin are from any cell source, in particular, yeast or mammalian cells (including human cells). The subunits may, for example, be calcineurin subunit A1 or A2. The CNI polypeptide may also be from any source (*e.g.*, yeast or human), and may be only a fragment of a complete CNI polypeptide (such as a c-terminal fragment). An exemplary c-terminal fragment of CNI is CN1c.

Further, included in the present invention, is a yeast cell carrying a mutation in the naturally-occurring copy of *CNI*, where the mutation prevents expression of a functional CNI protein from the genomic copy. Embodiments of this aspect of the present invention include deletion mutations within the coding region of the *CNI* gene, deletion of regulation regions of the *CNI* gene, and non-sense or mis-sense mutations in the *CNI* gene. Yeast cells having such mutations are useful, for example, in a method of identifying proteins of similar function to CNI. In one embodiment, a hybrid interaction screen is set up in a cell with a CNI deletion and a GAL4 protein binding domain-CNA fusion and a GAL4 activation domain-immunophilin fusion. Expression libraries are then screened to identify clones encoding proteins that potentiate an interaction of an immunophilin with calcineurin. This screen will identify CNI-coding sequences as well as other proteins with a similar function.



In a related embodiment, a yeast cell with a CNI deletion is used to identify CNI homologues (e.g., from other organisms, such as human) using a complementation assay or screen. Expression libraries (e.g., human lymphocyte expression libraries) are transformed into cells with a CNI deletion, and transformants are selected on their ability to complement the  
5 function of yeast CNI. An exemplary assay for selecting such transformants is exposure to hygromycin B. Cells which become more sensitive to hygromycin B following transformation are further analyzed to determine if the plasmid with which they were transformed contains an insert homologous to yeast CNI, or encoding a polypeptide with similar function to CNI.

The invention also includes a yeast cell carrying a mutation in the naturally-occurring  
10 genomic copy of a gene encoding calcineurin-interacting polypeptide, where the mutation prevents expression of a functional calcineurin-interacting polypeptide from the genomic copy. The mutation may be a null mutation, such as described in Example 8 below, or a different type of mutation, e.g., a nonsense or missense mutation. Nonsense and missense mutations may be generated using standard methods.

15 These and other objects and features of the invention will be more fully appreciated when the following detailed description of the invention is read in conjunction with the accompanying drawings.

#### **Brief Description of the Figures**

20 Figures 1A and 1B present schematic diagrams of sequences encoding the c-terminal portion of CNI (CN1c) fused to GAL-4 activation domain (GAL-4AD) (Fig. 1A), and sequences encoding CNI (Fig. 1B).

Figure 2A presents data from a  $\beta$ -galactosidase ( $\beta$ -gal) assay to detect the interaction of CN1c with the A1 subunit of calcineurin (CNA1), A2 subunit of calcineurin (CNA2), GAL-4  
25 binding domain (G4BD) and lamin C. A labeled schematic diagram corresponding to the data shown in Fig. 2A is presented in Fig. 2B to facilitate reference to individual groups of colonies.

Figure 3A presents data from a  $\beta$ -gal assay to detect the interaction of CN1c with CNA1 $\Delta$ C, CNA2 $\Delta$ C and CNB1. A labeled schematic diagram corresponding to the data  
30 shown in Fig. 3A is presented in Fig. 3B.

Figures 4A and 4C present data from  $\beta$ -gal assays to evaluate the effects of FK506 and the deletion of CNB1 on the interactions of CN1c with CNA and CNA $\Delta$ C. Labeled schematic diagrams corresponding to the data shown in Figs. 4A and 4C are presented in Figs. 4B and 4D, respectively.

Figures 5A and 5C present data from  $\beta$ -gal assays to evaluate the effects of FK506 and the overexpression of CNB1 on the interactions of CN1c with CNA and CNA $\Delta$ C. Labeled schematic diagrams corresponding to the data shown in Figs. 5A and 5C are presented in Figs. 5B and 5D, respectively.

- 5        Figures 6A, 6C and 6E present data from  $\beta$ -gal assays to evaluate the effects of FK506, cyclosporin A (CsA), and the deletion of CNB1 on the interaction of CN1c with CNA and CNA $\Delta$ C. Labeled schematic diagrams corresponding to the data shown in Figs. 6A, 6C and 6E are presented in Figs. 6B, 6D and 6F, respectively.

- 10       Figures 7A, 7C, 7E and 7G present data from  $\beta$ -gal assays to evaluate the effects of FK506, cyclosporin A (CsA), rapamycin and the overexpression of CNB1 on the interaction of CN1c with CNA and CNA $\Delta$ C. Labeled schematic diagrams corresponding to the data shown in Figs. 7A, 7C, 7E and 7G are presented in Figs. 7B, 7D, 7F and 7H, respectively.

- 15       Figure 8A presents data from a  $\beta$ -gal assay to evaluate the effects of overexpression of full-length CNI on FK506-dependent interaction of FKBP with CNA. A labeled schematic diagram corresponding to the data shown in Fig. 8A is presented in Fig. 8B.

Figures 9A and 9C present data from  $\beta$ -gal assays to evaluate the effects of overexpression of full-length CNI on FK506-dependent interaction of FKBP with CNB1. Labeled schematic diagrams corresponding to the data shown in Figs. 9A and 9C are presented in Figs. 9B and 9D, respectively.

- 20       Figure 10 presents an image of a protein blot of CN1c and CNA co-immunoprecipitate probed with anti-CNA2 antibody.

Figure 11 presents an image of a yeast RNA blot hybridized with a CN1c probe.

Figure 12 presents the DNA sequence of a 3.5 kb fragment of yeast chromosome 11 containing the coding sequence for a yeast CNI protein.

- 25       Figure 13 presents the amino acid sequence of a yeast CNI protein.

#### **Brief Description of the Sequences**

SEQ ID NO:1 presents the nucleotide sequence of a *Sau3AI* fragment containing the coding sequence for CN1c.

- 30       SEQ ID NO:2 presents the amino acid sequence of CN1c encoded by SEQ ID NO:1.

SEQ ID NO:3 presents the coding sequence presented in SEQ ID NO:1.

SEQ ID NO:4 presents the nucleotide sequence of a gene encoding a complete CNI protein.

SEQ ID NO:5 presents the amino acid sequence encoded by SEQ ID NO:4.

SEQ ID NO:6 presents the coding sequence presented in SEQ ID NO:4.

SEQ ID NO:7 presents the nucleotide sequence of PCR primer CNI-PCR-A.

SEQ ID NO:8 presents the nucleotide sequence of PCR primer CNI-PCR-B.

5     SEQ ID NO:9 presents the nucleotide sequence of a gene encoding the yeast CNA1 subunit of calcineurin.

SEQ ID NO:10 presents the amino acid sequence encoded by SEQ ID NO:9.

SEQ ID NO:11 presents the nucleotide sequence of a gene encoding the yeast CNA2 subunit of calcineurin.

SEQ ID NO:12 presents the amino acid sequence encoded by SEQ ID NO:11.

10     SEQ ID NO:13 presents the nucleotide sequence of a gene encoding the yeast CNB1 subunit of calcineurin.

SEQ ID NO:14 presents an amino acid sequence encoded by SEQ ID NO:13.

SEQ ID NO:15 presents an amino acid sequence encoded by SEQ ID NO:13.

SEQ ID NO:16 presents the coding sequence presented in SEQ ID NO:13.

15     SEQ ID NO:17 presents the amino acid sequence encoded by SEQ ID NO:16.

SEQ ID NO:18 presents a nucleotide sequence encoding CNA1Δc.

SEQ ID NO:19 presents the amino acid sequence encoded by SEQ ID NO:18.

SEQ ID NO:20 presents a nucleotide sequence encoding CNA2Δc.

SEQ ID NO:21 presents the amino acid sequence encoded by SEQ ID NO:20.

20     SEQ ID NO:22 presents the nucleotide sequence of PCR primer G4-PCR-A.

SEQ ID NO:23 presents the nucleotide sequence of PCR primer G4-PCR-B.

### **Detailed Description of the Invention**

#### **I. DEFINITIONS**

25     A "calcineurin-targeted immunosuppressant" is a compound that possesses *in vivo* immunosuppressive activity, and that interacts with an immunophilin to form a complex which is capable of inhibiting calcineurin.

"Interacting proteins" are proteins capable of specifically binding to one another, or associating with one another, in a cell or *in vitro*.

30     A calcineurin interacting (CNI) protein or polypeptide is a protein or polypeptide that is effective to enhance immunosuppressive effects of a calcineurin-targeted immunosuppressant by potentiating an interaction of an immunophilin with calcineurin. Preferably, a CNI protein or polypeptide is a protein or polypeptide having an amino acid sequence that is homologous to the sequence presented herein as SEQ ID NO:5.

"Substantially isolated" is used in several contexts and typically refers to the at least partial purification of a CNI protein or polypeptide fragment away from unrelated or contaminating components (*e.g.*, cytoplasmic contaminants and heterologous proteins). Methods and procedures for the isolation or purification of compounds or components of interest are described below (*e.g.*, affinity purification of fusion proteins and recombinant production of CNI polypeptides).

In the context of the present invention, the phrase "nucleic acid sequences," when referring to sequences which encode a protein, polypeptide, or peptide, is meant to include degenerative nucleic acid sequences which encode homologous protein, polypeptide or peptide sequences as well as the disclosed sequence.

Two nucleic acid fragments are considered to have "homologous" sequences if they are capable of hybridizing to one another (i) under typical hybridization and wash conditions, as described, for example, in Sambrook, *et al.*, pages 320-328, and 382-389, or (ii) using reduced stringency wash conditions that allow at most about 25-30% basepair mismatches, for example:  $2 \times$  SSC, 0.1% SDS, room temperature twice, 30 minutes each; then  $2 \times$  SSC, 0.1% SDS, 37°C once, 30 minutes; then  $2 \times$  SSC, room temperature twice, 10 minutes each. Preferably, homologous nucleic acid strands contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches. These degrees of homology can be selected by using wash conditions of appropriate stringency for identification of clones from gene libraries (or other sources of genetic material), as is well known in the art.

Two amino acid sequences or two nucleotide sequences (in an alternative definition for homology between two nucleotide sequences) are considered homologous (as this term is preferably used in this specification) if they have an alignment score of  $>5$  (in standard deviation units) using the program ALIGN, typically default mutation gap matrix and gap penalty (Dayhoff). The two sequences (or parts thereof) are more preferably homologous if their amino acids are greater than or equal to 40% using the ALIGN program mentioned above.

## II. OVERVIEW OF INVENTION

Experiments performed in support of the present invention demonstrate the identification and isolation of the nucleic acid sequence encoding a calcineurin interacting (CNI) protein. Further experiments performed in support of the present invention characterize the CNI protein, as well as a polypeptide containing only the c-terminal 306 amino acids of the CNI protein (CNIc). The experiments demonstrate that CNIc interacts specifically with the "A" subunits of calcineurin (CNA1 and CNA2), but not with calcineurin "B" subunit (CNB1).

The experiments also demonstrate that CN1c does not interact directly with FK506 binding protein (FKBP; Schreiber, *et al.*; with or without FK506), GAL4 binding domain (G4<sub>BD</sub>) or lamin C. The experiments also demonstrate that CN1c interacts with C-terminally truncated forms of CNA (CNAΔC), which have lost their autoinhibitory domains, though the interaction is somewhat weaker than with full length CNA proteins.

Additional experiments show that the interaction between CN1c and CNA is enhanced when CNB1 is deleted, and diminished when CNB1 is overexpressed, that the interaction between CN1c and CNA or CNAΔC is markedly enhanced by FK506 and by Cyclosporin A (CsA), but not rapamycin, and that overexpression of a full-length CNI protein enhances the interaction between CNA and FKBP (detectable only in the presence of FK506).

Additional experiments conducted in support of the present invention demonstrate that overexpression of the full-length CNI has no detectable effect on the interaction between CNB1 and CNA, and that in the presence of FK506 or CsA, overexpression of CNB1 no longer inhibits the interaction of CN1c with CNA.

It was also found that CNI deletion mutants are viable, both in wild-type and CN-deletion backgrounds, and that CNI deletion mutants in a CN-deletion background are more resistant to hygromycin B than normal CN-deletion mutants.

Co-immunoprecipitation experiments demonstrate that CN1c and CNA co-immunoprecipitate in the presence of FK506, and protein blot experiments show that CNI is expressed at low levels *in vivo*. RNA blot experiments show that CNI is encoded by a single message approximately 2.9 kb in length.

A comparison of the yeast CNI sequence with sequences present in nucleic acid and amino acid databases reveals no obvious homologous sequences have been identified in other organisms.

### III. CALCINEURIN

Experiments performed in support of the present invention were designed to identify polypeptides capable of interacting with calcineurin. Calcineurin (also called phosphoprotein phosphatase 2B or PP2B), has been characterized from many different tissues and organisms (Klee, *et al.*). It is a heterodimer of two subunits, of which the "A" subunit is about 61 kD in weight, possesses catalytic activity and also contains the association site for calmodulin. The "B" subunit contains four Ca<sup>2+</sup> binding sites and activates the A subunit. Calcineurin has little enzymatic activity, even in the presence of Ca<sup>2+</sup> and only becomes fully active when associated with calmodulin (Cyert).

Two A subunits (CNA1 and CNA2; Cyert, *et al.*, 1991) and one B subunit (CNB1, Cyert, *et al.*, 1992) have been cloned in yeast. Either CNA1 or CNA2 may associate with CNB1 to form a functional calcineurin heterodimer. Multiple isotypes of the A subunit have been cloned from a variety of organisms and are highly conserved (Klee, *et al.*). In particular, calcineurin subunits have been cloned from human tissue (see reviews by, for example, Klee, *et al.*, and Guerini, *et al.*).

#### IV. IMMUNOSUPPRESSANT DRUGS

FK506, cyclosporin A (CsA) and rapamycin, derived from fungi, inhibit the activation of T-cells by antigens. The compounds have proven highly effective at suppressing mammalian immune systems *in vivo*. In particular, CsA therapy in clinical settings has dramatically increased the success rate of transplantation therapy.

It is now known that FK506 and CsA exert their immunosuppressive effects, in part, by inhibiting the transcriptional activation of the interleukin-2 (IL-2) gene, whereas rapamycin appears to function by inhibiting the response of T-cells to IL-2, presumably by inhibiting a transduction pathway mediated by the IL-2 receptor.

The molecular mechanism of FK506 and CsA immunosuppressive action involves a group of small, abundant intracellular proteins termed immunophilins, which bind with a high affinity to the immunosuppressants (Schreiber). At least two classes of immunophilins are known to exist. One class, termed cyclophilins, binds to CsA, while another class, the FK506-binding proteins (FKBPs) binds FK506 and rapamycin. Many immunophilin genes, from a variety of organisms, have been cloned, and appear to be highly conserved from simple eukaryotes to mammals.

It is believed that FK506 and CsA-induced immunosuppression is due to the binding of complexes, formed by binding of immunosuppressants FK506 and CsA bound to one of their respective immunophilins, to the catalytic subunit of calcineurin (Schreiber, *et al.*, Liu, *et al.*, Foor, *et al.*, Weiss, *et al.*). The binding of such a complex to an (A) subunit inhibits activation of calcineurin by increased intracellular calcium, which in turn prevents calcineurin from activating transcription factor NF-AT. Since IL-2 is one of the genes controlled by NF-AT in T-cells, inhibition of the transcription factor inhibits the production of IL-2, resulting in immunosuppression (Clipstone, *et al.*).

FK506 and CsA are widely used in organ transplantation to prevent host rejection. However, both drugs are known to have many undesired side-effects such as neurotoxicity, nephrotoxicity, hypertension, and metabolic disorder. Accordingly formulations effective to

increase a target cell's sensitivity to these drugs may be useful in alleviating some of the aforementioned side-effects. Specifically, CNI and its homologues or derivatives, administered at appropriate levels, may be able to increase the sensitivity of CN to FK506/CsA and reduce the necessary dosage thus reducing or eliminating the side-effects of these drugs.

5

#### V. TWO HYBRID PROTEIN INTERACTION ASSAYS

Two hybrid protein interaction assay methods (two hybrid protein-protein interaction screens) provide a simple and sensitive means to detect the interaction between two proteins in living cells. The assays are based on the finding that most eukaryotic transcription activators are modular (e.g. Brent, *et al.*), *i.e.*, that the activators typically contain activation domains that activate transcription, and DNA binding domains that localize the activator to the appropriate region of a DNA molecule.

The development of two hybrid protein interaction assays was made possible by the observation that the DNA binding domain does not need to be physically located on the same polypeptide as the activation domain (Ma, *et al.*, Triezenberg, *et al.*), raising the possibility that transcription of reporter genes could be used as an assay to detect protein interactions.

The utility of two hybrid systems for detecting interactions between two interacting proteins was fully realized by the observation that protein interactions could be detected if two potentially-interacting proteins were expressed as fusions, or chimeras (Fields, *et al.*). A first fusion protein contains one of a pair of interacting proteins fused to a DNA binding domain, and a second fusion protein contains the other of a pair of interacting proteins fused to a transcription activation domain. The two fusion proteins are independently expressed in the same cell, and interaction between the "interacting protein" portions of the fusions reconstitute the function of the transcription activation factor, which is detected by activation of transcription of a reporter gene.

At least two different cell-based two hybrid protein-protein interaction assay systems have been used to assess binding interactions and/or to identify interacting proteins. Both employ a pair of fusion hybrid proteins, where one of the pair contains a first of two "interacting" proteins fused to a transcription activation domain of a transcription activating factor, and the other of the pair contains a second of two "interacting" proteins fused to a DNA binding domain of a transcription activating factor.

The yeast GAL4 two hybrid system (Fields, *et al.*, Chien, *et al.*, Durfee, *et al.*, Bartel, *et al.*), utilized for experiments performed in support of the present invention, was developed

to detect protein-protein interaction based on the reconstitution of function of GAL4, a transcriptional activator from yeast, by activation of a *GAL1-lacZ* reporter gene.

Like several other transcription activating factors, GAL4 contains two distinct domains, a DNA binding domain and a transcription activation domain. Each domain can be  
5 independently expressed as a portion of a fusion protein composed of the domain, and a second, "bait" interacting protein. The two fusion proteins are then independently expressed together in a cell. When the two GAL4 domains are brought together by a binding interaction between the two "interacting" proteins, transcription of a reporter gene under the transcriptional control of GAL4 is initiated. The reporter gene typically has a promoter containing GAL4  
10 protein binding sites (GAL upstream activating sequences, UAS<sub>G</sub>). Exemplary reporter genes are the *GAL1-lacZ*, and *GAL1-HIS3* reporter genes used in experiments described herein.

A second two hybrid system, described in detail in Ausubel, *et al.*, utilizes a native *E. coli* LexA repressor protein, which binds tightly to appropriate operators. A plasmid is used to express one of a pair of interacting proteins (the "bait" protein) as a fusion to LexA. The  
15 plasmid expressing the LexA-fused bait protein is used to transform a reporter strain of yeast, such as EGY48, that contains pSH18-34.

In this strain, binding sites for LexA are located upstream of two reporter genes. In the first reporter system, the upstream activation sequences of the chromosomal LEU2 gene—required in the biosynthetic pathway for leucine (Leu)—are replaced in EGY48 with *lexA*  
20 operators, permitting selection for viability when cells are plated on medium lacking Leu. In the second reporter system, EGY48 harbors a plasmid, pSH18-34, that contains a *lexA* operator-*lacZ* fusion gene, permitting discrimination based on color when the yeast is grown on medium containing Xgal (Ausubel, *et al.*).

To screen a library with the LexA system, the library uses the inducible yeast GAL1  
25 promoter to express proteins as fusions to an acidic domain ("acid blob") that functions as a portable transcriptional activation motif ("act"), and to other useful moieties. Expression of library-encoded proteins is induced by plating transformants on medium containing galactose (Gal), so yeast cells containing library proteins that do not interact specifically with the bait protein fail to grow in the absence of Leu. Yeast cells containing library proteins that interact  
30 with the bait protein form colonies within 2 to 5 days, and the colonies turn blue when the cells are streaked on medium containing Xgal. The plasmids are isolated and characterized by a series of tests to confirm specificity of the interaction with the initial bait protein. Those found to be specific are ready for further analysis (e.g., sequencing).



LexA and GAL4 each have different properties that should be considered when selecting a system. LexA is derived from a heterologous organism, has no known effect on the growth of yeast, possesses no residual transcriptional activity, can be used in GAL4<sup>+</sup> yeast, and can be used with a Gal-inducible promoter. Because GAL4 is an important yeast  
5 transcriptional activator, experiments must be performed in *gal4* yeast strains to avoid background from endogenous GAL4 activating the reporter system.

Both two hybrid systems have been successfully used for isolating genes encoding proteins that bind a target protein and as simple protein binding assays (*e.g.*, Yang, *et al.*, Gyuris, *et al.*), and both can be applied to methods of the present invention.

10 Both gene isolation and protein binding assay applications of the GAL4 system are described in Examples below.

#### VI. SPECIFIC EMBODIMENTS

Example 1 demonstrates application of an exemplary two hybrid protein-protein  
15 interaction screen (Materials and Methods, section D) to the screening of three pGAD yeast fusion libraries, carrying fusions between the transcription activating domain of yeast protein GAL4 (G4AD) and yeast genomic DNA *Sau3AI* fragments, in all three reading frames. The libraries are screened to identify polypeptides, encoded by the *Sau3AI* fragments, capable of interacting with catalytic (A) subunits of calcineurin, expressed as fusions with the GAL4  
20 protein binding domain (GBT-CNA fusions).

Three sets of yeast cells harboring pGBT-CNA1 *TRP1* (GBT-A1) hybrid plasmid and a GAL4-activated LacZ reporter gene are each transformed with one of the three reading-frame libraries. Construction of the plasmids used is described in Materials and Methods, sections B and C. Cells transformed with a plasmid encoding a protein fusion capable of interacting  
25 with the CNA subunit fusion are selected using a  $\beta$ -galactosidase ( $\beta$ -gal) assay on plates containing the chromogenic substrate X-gal (Materials and Methods, section E). Results of the  $\beta$ -gal assay are confirmed using a growth assay (Materials and Methods, section F). False positives are eliminated by colony purification (re-streaking for single colonies), PCR experiments using GAL4 primers, and testing against a number of test fusions by  $\beta$ -gal assays  
30 on transformed haploid or mated diploid reporter strains.

A yeast clone encoding a polypeptide capable of specifically interacting with CNA polypeptide fusions is identified and sequenced. The sequence of the *Sau3AI* fragment is presented as SEQ ID NO:1. The coding sequence forming the open reading frame is presented as SEQ ID NO:3. The polypeptide encoded by the open reading frame is presented as SEQ

ID NO:2. The open reading frame encodes 306 amino acids followed by a stop codon in frame with the coding sequence of GAL4, but does not contain an in-frame ATG start site upstream of the stop codon. This is consistent with the fragment encoding a c-terminal portion of a complete gene product. The clone is termed CN1c, with the lowercase "c" representing "c-terminal".

Figure 1A shows a schematic representation of the nucleic acid sequence encoding the GAL4AD-CN1c fusion protein. The stippled portion between GAL4AD and CN1c represents a linker discussed in the Materials and Methods section, as well as in Example 1. Also indicated in the Figure are the approximate locations of in-frame start (ATG) and stop codons, and *Sau3A*I restriction sites.

Example 1 further describes the identification of a  $\lambda$  clone encoding a full length sequence version of CN1c, termed CN1. The polypeptide encoded by the sequence is termed CN1 protein. The clone is identified by hybridization screening of a panel of  $\lambda$  clones spanning the yeast genome using a 1.22 kb  $^{32}$ P-labeled probe generated from CN1c.

Phage lysates of the  $\lambda$  clone are amplified, purified, restriction-mapped and used as a DNA source for subsequent cloning experiments. A 3.16 kb *Mun*I/*Eco*RV fragment from the  $\lambda$  clone insert contains the coding sequence of CN1. The sequence of the 3.16 kb *Mun*I/*Eco*RV fragment is encompassed by the 3.5 kb sequence presented as SEQ ID NO:4, and a schematic diagram of the sequence is shown in Figure 1B. This sequence contains the entire 2.75 kb coding sequence of CN1 (SEQ ID NO:6; graphically indicated in Figure 1B by the locations of the "ATG" and "Stop" codons).

A search of known DNA and protein sequences turns up no obvious matches or homologies to genes in other organisms. Accordingly, CN1 may represent a new type of calcineurin interacting protein.

The methods referred to above may also be applied to the screening of, for example, a human cDNA library using an appropriate two-hybrid protein interaction screen. The "bait" protein in the interaction screen (*e.g.*, the protein analogous to CNA1 in Example 1) may be of yeast origin (*e.g.*, CNA1), but is preferably of human origin (*e.g.*, a human calcineurin "A" subunit; Klee, *et al.*). The bait protein is expressed in the cell (*e.g.*, a yeast cell) used for the two hybrid interaction screen as a fusion to a domain of a transcription activating factor (*e.g.*, the DNA binding domain of GAL4). The library may be a human DNA library in a vector (*e.g.*, pGAD) effective to express library sequences as fusions to a complimentary domain of the transcription activating factor (*e.g.*, the activation domain of GAL4). Libraries of human sequences can be derived from a number of sources including genomic DNA, such as yeast

artificial chromosome (YAC) constructs carrying genomic human DNA, or cDNA generated from a variety of cell types (*e.g.*, activated T-cells).

Example 2 details a  $\beta$ -gal assay to determine the specificity of binding of CN1c to subunits of calcineurin. Exemplary results are shown in Figure 2A. The legend for Fig. 2A is shown in Figure 2B. Numbers in the legend refer to locations of yeast colonies expressing particular combinations of plasmid constructs (indicated in Example 2).

A comparison of the intensities of the blue  $\beta$ -gal reaction product indicates that CN1c interacts strongly with CNA1 (21, 22 and 23), and somewhat less strongly with CNA2 (24). Neither cells containing AS-lamin with GAD-CN1c (25), nor cells containing only GBT with GAD-CN1c (26) show a detectable signal above background. Two subunits of calcineurin (CNA1, CNB1) known to interact with each other are used as a positive control for the assay (20). The data presented in Figure 2A show that CN1c interacts specifically with CNA1 and CNA2, but not with G4BD or lamin C.

A similar set of experiments, illustrated in Fig. 3A, is conducted using constitutively-active CNA subunits, as well as calcineurin subunit CNB1. As described in the Materials and Methods section below, CNA1 $\Delta$ C and CNA2 $\Delta$ C are each missing a C-terminal portion of the protein containing an autoinhibitory domain. Exemplary results from these experiments are shown in Figure 3A. The legend for Fig. 3A is shown in Figure 3B. The data show that GBT-A1 $\Delta$ C and GAD-CN1c (28) gives a definite positive signal, while GBT-A2 $\Delta$ C and GAD-CN1c (29) is weaker, though still detectable above its background (*i.e.* GBT-A2 $\Delta$ C and GAD; 31). The signal from GBT-B1 and GAD-CN1c (32) is not detectable above vector background (GBT-B1 and GAD; 33). Positive controls GBT-A1 $\Delta$ C and GAD-B1 (34) and GBT-A2 $\Delta$ C and GAD-B1 (35) give strong signals. The data presented in Figure 3A show that CN1c interacts specifically with CNA1 $\Delta$ C and CNA2 $\Delta$ C, but not with CNB1.

Example 3 details the effects of immunosuppressant drugs on binding of CN1c to calcineurin in B1<sup>+</sup>, B1 Deletion and B1 Overproducing Yeast Strains. The yeast strains are assayed for  $\beta$ -gal activity as above to determine if the immunosuppressant drugs FK506, cyclosporin A (CsA) and rapamycin affect the binding of CN1c to subunits of calcineurin. The experiments are performed in yeast strains wild-type for the CNB1 subunit, null for the CNB1 subunit, and in yeast transformed with a high efficiency expression vector containing DNA encoding the CNB1 subunit.

Exemplary data are shown in Figures 4A, 4C, 5A and 5C. The legends for Figs. 4A and 4C are shown in Figures 4B and 4D, respectively. Figs. 4A and 5A illustrate experiment

performed without FK506, while experiments shown in Figs. 4C and 5C were performed with FK506.

The interactions of various combinations of proteins expressed by constructs indicated in Example 3 was studied in three yeast strains, one of which is null for the CNB1 subunit of calcineurin (Y153b; at 36-40), while the others (Y190; at 41-46 and Y526 at 47-51) are wild-type for CNA1, CNA2 and CNB1.

The data, shown in Fig. 4A (no added drugs), illustrate that deleting the endogenous host CNB1 gene potentiates, or enhances, interactions between CN1c and calcineurin subunits CNA1, CNA2, CNA1ΔC and CNA2ΔC. Comparison of corresponding colonies in Figs. 4A and 4C shows the effects of FK506 on CN1c-CNA/CNAΔC interactions. The drug enhances interactions under all except control conditions. The effect is most striking in yeast strains wild-type for the CNB1 subunit (*e.g.*, compare 50a with 50b, and 51a with 51b).

The drug also markedly enhances, or potentiates CN1c-CNA/CNAΔC interactions under conditions where the CNB1 subunit is overexpressed. Figure 5A shows the effect of overexpressing CNB1 on CN1c-CNA/CNAΔC interactions in the absence of drug. Colonies expressing B1/YEp352 (53a, 55a-58a) have reduced signal as compared with controls expressing only YEp352 (52a, 54a). Inclusion of FK506 in the plating medium (Fig. 5C), however, enhances, or potentiates interactions in all colonies, except the negative controls (56).

The data presented in Figures 4A, 4C, 5A and 5C demonstrate that the interaction of CN1c with CNA and CNAΔC is markedly enhanced by FK506. The interaction is also enhanced by deletion of CNB1 and diminished by overexpression of CNB1, and the inhibitory effect of CNB1 overproduction is overcome by the stimulatory effect of FK506.

Stated another way, inclusion of a small molecule immunosuppressant (FK506) potentiates an interaction between two fusion hybrid proteins, where one of the two proteins contains an (A) subunit of calcineurin, and the other protein contains a CNI polypeptide. The potentiation is particularly strong when the cell is further modified to cause overexpression of a "B" subunit of calcineurin by said cell (*e.g.*, the expresses B1/YEp352).

In the present case, a yeast cell is modified to cause overexpression of a "B" subunit of calcineurin (CNB1) by transforming the cell with B1/YEp352 (construction described below). A cell may be modified to cause overexpression of a "B" subunit of calcineurin in other ways as well, such as, for example, transformation with other types of expression vectors encoding a "B" subunit of calcineurin, or treatment with a substance that upregulates a promoter controlling expression of an endogenous (B) subunit of calcineurin.

In light of the effects of FK506 on CN1c-CNA/CNAΔC interactions, two other immunosuppressants, cyclosporin A and rapamycin, were examined in similar experiments. The results of these experiments are shown in Figures 6A-6F and 7A-7H.

5 The data presented in Figures 6A and 6C are essentially equivalent to those presented in Figures 4A and 4C, respectively. Data shown in Figure 6E demonstrate that like FK506, the immunosuppressant cyclosporin A is also effective to enhance interaction of CN1c with CNA and CNAΔC. Both FK506 and cyclosporin A are known to exert their immunosuppressive effects through calcineurin (Cyert).

10 Similarly, data shown in Figures 7A and 7C support data in Figures 5A and 5C, and further, demonstrate that there is no detectable interaction between FK506 binding protein (FKBP) and CN1c. Results shown in Fig. 7E demonstrate that cyclosporin A has a similar effect to FK506 in cells overexpressing CNB1 — that is, it enhances the interactions between CN1c and CNA/CNAΔC.

15 In contrast, data presented in Fig. 7G show that the immunosuppressant rapamycin has no detectable effect on CN1c-CNA/CNAΔC interactions (compare Fig. 7G with Fig. 7A).

Taken together, the data presented in Figs. 6A, 6C, 6E 7A, 7C, 7E and 7G show that like FK506, cyclosporin A (CsA), but not rapamycin, enhances the interaction of CN1c with CNA and CNAΔC, and that CN1c doesn't interact with FKBP with or without FK506.

20 Example 4 describes experiments to assess effects of CN1c on FKBP/FK506 binding to calcineurin. Figure 8A presents exemplary data from studies to assess the effect of CN1 on FKBP-mediated FK506 interactions with CNA2. The legend for Fig. 8A is shown in Figure 8B. Data in Figure 8A demonstrate that, in the absence of FK506, FKBP and CNA2 show no detectable interaction (84). In the presence of FK506, however, the proteins interact (85), presumably because FK506 forms a complex with FKBP, which then binds CNA2 (Cyert).

25 Data in Figure 8A further show that, in the absence of FK506, CN1 has no effect on the lack of interaction between FKBP and CNA2 (86). In the presence of FK506 (87), however, CN1 potentiates, or enhances the binding between FKBP and CNA2 (compare the intensity at 87 with that at 85). This effect suggests that CN1 and similar compounds may be employed to increase the sensitivity of calcineurin to immunosuppressant drugs that act on it, and in this way, decrease the amount of the immunosuppressant required for a particular level  
30 of immunosuppression.

Experiments illustrated in Figures 9A and 9B demonstrate that FK506 has little or no effect on the binding of CNA to CNB1. The legends for Fig. 9A and 9B are shown in Figures 9B and 9D, respectively. The data in Fig. 9A show that overexpression of the full-length CN1

clone markedly enhances the FK506-dependent interaction of FKBP with CNA, although it doesn't affect the interaction between CNA and CNB1.

Example 5 presents co-immunoprecipitation of CN1c (carrying an HA epitope tag) and CNA. Immunoprecipitation is carried out with anti- HA monoclonal antibody and the immune complex, resolved by SDS-PAGE, is detected with anti-CNA2 polyclonal antibody and visualized with goat anti-rabbit antibody using the "ECL" method (Amersham, Arlington Heights, IL). The results, shown in Figure 10, demonstrate that CN1c is capable of binding to CNA2 tightly enough for the complex to be co-immunoprecipitated. Similar methods may be employed to isolate a CNI analog from other cell sources, including mammalian (specifically human).

Example 6 describes yeast RNA blots hybridized with a CN1c probe. Exemplary data are shown in Figure 11. A single message of approximately 2.9 kb is detected. The data indicate that CNI is an expressed gene encoding a 2.9 kb message in yeast.

Example 8 details the construction of cni null mutants. The null mutants are employed to assess if *CNI* is required for viability in yeast, and to test hygromycin B sensitivity. The experiments indicate that CNI is not essential for viability, since CNI deletion mutant strains can survive, but that CNI deletions render a host more resistant to hygromycin B. The effect is particularly pronounced in both MCY300-1 (*cna1<sup>-</sup>cna2<sup>-</sup>*) and DD12 (*cnb1<sup>-</sup>*), suggesting that CNI functions as a suppressor of CN mutant's sensitivity to hygromycin B.

It will be understood that all of the above methods and experimental manipulations are amenable to being done with interacting polypeptides from organisms other than yeast. In particular, calcineurin subunits, CNI polypeptides, immunophilins and the like may be of mammalian origin, *e.g.*, human origin.

## VII. UTILITY

Methods and compositions of the present invention may be applied in a number of different ways. Following the guidance presented herein, one of skill in the art may isolate nucleic acids encoding additional CNI polypeptides, for example, a human CNI polypeptide.

In one approach, a yeast strain carrying a mutation of the *CNI* gene, *e.g.*, a deletion, is used to clone heterologous sequences (*e.g.*, human sequences) by complementation. A library of genomic DNA or, preferably, cDNA from an organism (*e.g.*, human) and tissue (*e.g.*, lymphocyte cells) of choice is cloned into a vector that can be maintained in yeast. Preferably, the vector contains a yeast promoter effective to express the heterologous sequences in yeast cells. Several heterologous libraries suitable for expression in *Saccharomyces*

*cerevisiae* containing DNA from *S. pombe* (Beach, *et al.*) and *Drosophila* have been constructed.

The library is transformed into a suitable yeast strain carrying a *cni* mutation, and transformants are selected using a suitable complementation assay. For example, transformants  
5 may be screened for increased hygromycin sensitivity, as experiments described herein indicate that *cni* deletion mutants possess a decreased sensitivity to hygromycin B (Example 8). The screen may be made more effective by using a yeast strain that is hypersensitive to hygromycin B, such as a strain deficient for a subunit of calcineurin (Example 8).

Alternatively, human CNI DNA sequences may be isolated by directly screening a  
10 library, *e.g.*, a lymphocyte cDNA library, for clones hybridizing with a yeast CNI nucleic acid probe. The generation of an exemplary yeast CNI nucleic acid probe is described in Example 1.

In another approach, particularly advantageous for isolating sequences expressed at low levels, a CNI nucleic acid probe may be used to screen a genomic library, *e.g.*, a human  
15 genomic library, to isolate a sequence that may be used to design probes or primers that may match the target sequence better than the yeast sequence. Such primers may be used with, for example, PCR, to isolate longer fragments from a tissue-specific library.

In yet another approach, an antibody generated against CNI polypeptide is used to immunoprecipitate a CNI polypeptide from an organism and/or tissue of choice. The protein  
20 may then be micro-sequenced, and the sequence utilized to design degenerate primers useful for isolating a cDNA.

CNI polypeptides of the present invention, particularly CNI fragments that retain a desired binding activity, may be used as lead compounds useful for the development of small molecules having cellular functions similar to those of the CNI-polypeptides, that is, molecules  
25 effective to enhance immunosuppressive effects of calcineurin-targeted immunosuppressants by potentiating an interaction of an immunophilin with calcineurin.

CNI-polypeptides of the present invention may also be employed in a method of increasing sensitivity of cells to calcineurin-affecting immunosuppressant drugs. In this method, a CNI-polypeptide is introduced into the cell typically prior to or at the same time as  
30 contacting the cell with an immunosuppressant drug, such as FK506. The polypeptide may be delivered by any suitable means effective to deliver polypeptides to selected cells.

Alternatively, nucleic acids encoding CNI polypeptides may be used in appropriate expression vectors as a genetic therapy tool to potentiate the immunosuppressive effects of

calcineurin-targeting immunosuppressant drugs. The vectors may be targeted to selected cells, such as T-cells, to increase their sensitivity to a given systemic dose of an immunosuppressant.

Another utility of the present invention includes methods of screening for substances that up-regulate expression of CNI polypeptides, *i.e.*, substances that affect transcription. Such substances are useful for sensitizing cells to immunosuppressant drugs. In this method, the CNI promoter can be attached to a gene that functions as a selectable marker (for use in genetic selections to screen test substances) or to a reporter gene (for use in evaluating the effect on CNI transcription by test substances).

In another aspect of the present invention, the CNI-polypeptides, for example, mammalian homologue polypeptides of CNI, have potential use as therapeutic agents for both human and veterinary use. For example, CNI-polypeptides may be used in a method of enhancing immunosuppression in a test subject. In this method, the CNI-polypeptide is administered to the subject in a pharmaceutically-acceptable formulation and at a concentration effective to potentiate the interaction of an immunosuppressant/immunophilin complex with a subunit of calcineurin. The method may also include contacting the CNI-polypeptide with a cell under conditions effective to permit uptake of the protein into the cell in order to increase sensitivity of the cell to immunosuppressants. A CNI polypeptide used in such methods may be modified to be more suitable for administration or to be more effective in a cell. For example, a CNI polypeptide may be modified to eliminate PEST motifs, which are typically found in proteins with short half-lives, to extend the effective lifetime of the polypeptide in the target cell.

The following examples illustrate, but in no way are intended to limit the present invention.

25

#### MATERIALS AND METHODS

Unless indicated otherwise, chemicals and reagents were obtained from Sigma Chemical Company (St. Louis, MO) or Mallinckrodt Specialty Chemicals (Chesterfield, MO), restriction endonucleases were obtained from New England Biolabs (Beverly, MA), and other modifying enzymes and biochemicals were obtained from Pharmacia Biotech (Piscataway, NJ) or Boehringer Mannheim (Indianapolis, IN). FK506 was obtained from Fujisawa USA, Inc. (Deerfield, IL), cyclosporin A was obtained from Sandoz (Basel, Switzerland), and rapamycin was obtained from Wyeth-Ayerst (Princeton, NJ). Materials for media for yeast growth and culture were obtained from DIFCO (Detroit, MI). Unless otherwise indicated, manipulations



of yeast, bacteria, nucleic acids, proteins and antibodies were performed using standard methods and protocols (*e.g.*, Guthrie, *et al.*, Sambrook, *et al.*, Ausubel, *et al.*, Harlow, *et al.*, and Rose, *et al.*).

5           A.     Buffers

Z buffer: 60 mM Na<sub>2</sub>HPO<sub>4</sub>-7H<sub>2</sub>O, 40 mM NaH<sub>2</sub>PO<sub>4</sub>-H<sub>2</sub>O, 10 mM KCl, 1 mM MgSO<sub>4</sub>-7H<sub>2</sub>O and 50 mM  $\beta$ -mercaptoethanol (pH 7.0).

          B.     Plasmids, Libraries and Yeast Strains

10           Plasmids pGBT9 (GBT), carrying GAL4 DNA-binding domain (amino acid residues 1-147; G4BD) and TRP1, and pGAD (GAD), carrying GAL4 activation domain (amino acid residues 768-881; G4AD) and LEU2; three pGAD libraries carrying fusions between G4AD and yeast genomic *Sau3AI* partial-digest fragments in each frame; and the yeast *GAL1-lacZ* reporter strain SFY526 (Y526; *MATa ura3-52 ade2-101 lys2-801 his3-200 trp1-901 leu2-3,112*  
15 *can1 gal4-542 gal80-538 URA3::GAL1-lacZ*) were obtained from Stanley Fields (State University of New York at Stony Brook, Stony Brook, NY; Chien, *et al.*, Bartel, *et al.*). The libraries were constructed with linkers between the GAL4 activation domain and the *Sau3AI* fragments. The sequences of the linkers were 5'-ATCG-3' for the first library, 5'-ATCCG-3' for the second library, and 5'-ATCCCG-3' for the third library. In this way, the yeast  
20 genomic *Sau3AI* fragments were cloned in all three reading frames relative to G4AD.

          Plasmids pAS2 (AS) carrying G4BD and TRP1, and pAS-lamin (AS-lamin) containing a sequence encoding a G4BD-lamin C fusion; and yeast reporter strains Y190 (*MATa ura3-52 ade2-101 his3- $\Delta$ 200 trp1-901 leu2-3,112 cyh2 $\Delta$ , gal4 $\Delta$  gal80 $\Delta$  URA3::GAL-lacZ LYS2::GAL-HIS3*), a derivative of Y153 carrying dual indicator genes (GAL-lacZ and GAL-  
25 HIS3), and Y187 (*MATa ura3-52 ade2-101 lys2-801 his3-200 trp1-901 leu2-3,112 gal4 $\Delta$  gal80 $\Delta$  URA3::GAL-lacZ*) carrying *GAL-lacZ* reporter were obtained from Stephen Elledge (Baylor College of Medicine, Houston, TX; Durfee, *et al.*). Yeast strain Y153b1 (*cnb1::ADE2*) was derived from Y153.

*E coli* strain JBe181 (*leuB600 trpC9830*) was obtained from Ira Herskowitz (University  
30 of California at San Francisco, San Francisco, CA). Protease-deficient yeast strain BJ2407 (Guthrie, *et al.*) was obtained from the Yeast Genetic Center (University of California at Berkeley, Berkeley, CA).

### C. GAL4-Calcineurin Fusions

GAL4-calcineurin (GAL4-CN) fusions, GBT-A1 (G4BD-CNA1), GBT-A2 (G4BD-CNA2), GBT-B1 (G4BD-CNB1), GAD-A1 (G4AD-CNA1), GAD-A2 (G4AD-CNA2), and GAD-B1 (G4AD-CNB1) were constructed as follows. Plasmids containing inserts encoding  
5 CN subunits CNA1 (SEQ ID NO:9; Cyert, *et al.*, 1991), CNA2 (SEQ ID NO:11; Cyert, *et al.*, 1991) and CNB1 (SEQ ID NO:13; Cyert, *et al.*, 1992) were subjected to site-directed mutagenesis (Kunkle) to introduce a *Bam*HI site just upstream of each subunit's initiation codon in the second reading frame. DNA prepared from the mutated plasmids was digested with *Bam*HI and *Xho*I, and the resulting *Bam*HI-*Xho*I fragments, each containing a full-length coding  
10 sequence, were cloned into GBT or GAD that had been cut with *Bam*HI and *Sal* I. The resulting plasmids encoded in-frame fusions of the CN subunits with G4BD or G4AD.

Plasmids encoding CNA protein variants with truncated C-termini (GBT-A1ΔC, GBT-A2ΔC) were constructed by introducing stop codons after amino acid residues 509 (CNA1) and 502 (CNA2). The 44-residue deletion in GBT-A1ΔC removed the autoinhibitory domain of  
15 CNA1, while the 102-residue deletion in GBT-A2ΔC removed both the autoinhibitory and the calmodulin-binding domains of CNA2 (Cyert, *et al.*, 1991).

Plasmid GBT-FKBP, containing an FK506 binding protein (FKBP) gene fused to the GAL4 binding domain, was constructed by introducing a *Bgl*II site upstream of the initiation codon and a *Bam*HI site downstream of the stop codon of FKBP12 (Heitman, *et al.*) and  
20 ligating the *Bgl*II-*Bam*HI fragment into GBT cut with *Bam*HI.

Plasmid B1/YEp352 was constructed to contain the full coding sequence of CNB1 (SEQ ID NO:13, Cyert, *et al.*, 1992) as a 1.4 kb *Bam*HI-*Eco*RI fragment encompassing the sequence presented as SEQ ID NO:13 (812 bp; contains the coding sequence), in the multicopy plasmid YEp352(HIS), which is derived from YEp352 (URA) (Hill, *et al.*).

25 Plasmid CNI/YEp352(HIS) (also referred to as CNIH) was constructed by ligating a 3.16 kb *Mun*I-*Eco*RV fragment, containing the full coding sequence of CNI, from plasmid CNI7.1 (construction described below) into YEp352(HIS) cut with *Eco*RI and *Sma*I. Plasmids CNI/YEp352(TRP) (also referred to as CNIT) and CNI/YEp352(URA) (also referred to as CNIU) were similarly constructed using the 3.16 kb fragment and YEp352(TRP) or  
30 YEp352(URA), respectively (Hill, *et al.*).

Plasmids A1/YEp351 and A2/YEp352 were constructed to contain the full coding sequences of CNA1 (SEQ ID NO:9, Cyert, *et al.*, 1991) in YEp351 (Hill, *et al.*) and CNA2 (SEQ ID NO:11, Cyert, *et al.*, 1991) in YEp351 and YEp352, respectively. A1/YEp351 was constructed by ligating a 2.9 kb *Sac*I-*Hind*III fragment from clone CNA1 (Cyert, *et al.*, 1991)

into YEp351(HIS) cut with *SacI* and *HindIII*. A2/YEp352 was similarly constructed by ligating a 3 kb *SpeI-HindIII* fragment from clone CNA2 (Cyert, *et al.*, 1991) into YEp352(HIS) cut with *XbaI* and *HindIII*.

5 All GAL4-CN fusions were verified by DNA sequencing (Sanger, *et al.*) using "SEQUENASE 2.0" sequencing kits (United States Biochemical, Cleveland, OH), and were subjected to the following tests. The functionality of the fusion proteins was assayed by determining whether they could complement the appropriate *cn<sup>-</sup>* mutant phenotypes, using assays to measure the sensitivity to pheromone and  $Mn^{2+}$  (Reneke, *et al.*, Cyert, *et al.*, 1991). All of the GAL4-CN fusions were functional in this assay.

10 The fusion proteins were also tested for their ability to activate the reporter gene in the absence of the complementary GAL-4 domain fusion (*i.e.*, in the presence of the complementary GAL4 domain not fused to a second protein, for example, G4BD-A1 vs G4AD) using the two-hybrid interaction assay described below. Only GBT-B1 and GAD-A1 were able to activate the reporter gene at low levels without the complimentary GAL-4 domain fusion --  
15 assays with the other fusion proteins in the absence of the complimentary GAL-4 domain fusion showed no detectable levels of expression.

The two-hybrid interaction assay was also used to test the ability of the fusions to interact specifically with another fusion containing complimentary GAL4 and CN domains (*e.g.*, G4BD-A1 interacting with G4AD-B1). All CN hybrids were able to react specifically  
20 and result in an activation of the reporter gene that was clearly detectable above background. The high specificity witnessed in these experiments indicates that the GAL4 two-hybrid system can reliably be used to assay interactions between CN and other proteins.

#### D. Yeast GAL4 Two-Hybrid System for Detecting Protein-Protein Interaction

25 In the library screen, described in more detail in Example 1A, the yeast strain Y190, harboring the hybrid plasmid carrying the GAL4 binding domain fused to the A1 subunit of calcineurin (G4BD-CNA1), was transformed with fusion libraries carrying yeast genomic DNA *Sau3AI* fragments fused to the GAL4 activation domain. Transformants that were able to express the reporter genes, *i.e.*, able to grow on -His + 3-AT and to score blue in  $\beta$ -gal assay,  
30 were selected as candidate positives. These candidate positives potentially contain library DNA fragments encoding proteins that physically interact with CNA1.

In another application described herein, the two-hybrid system was used to test for interactions between CNA (fused to one of the GAL4 domains) and CNB1 (fused to the other GAL4 domain), and between CNA and FKBP. Additional experiments tested a clone, CN1c,

isolated using the library screen, against a series of proteins fused to the complementary GAL4 domain under various conditions to test whether CN1c interacts with CNA subunits, and if so, how the interactions are affected by various conditions.

5           E.     Color Development ( $\beta$ -gal) Assay

Yeast reporters harboring both G4BD and G4AD fusions (and a third non-fusion plasmid in some cases) were monitored for  $\beta$ -gal activity as follows. Purified yeast transformants were patched onto selective plates with or without other test reagents. After growing 3 days at 30°C, colonies were lifted onto nitrocellulose filters, permeabilized in liquid  
10 nitrogen as above, placed on Whatman No. 1 paper in petri dishes containing 0.1% X-Gal in Z buffer (see above), and incubated at 30°C for 12 hours. Blue color begins to appear in positive colonies between about one half and ten hours into the incubation period.

Exemplary images obtained using the color development assay are presented in Figures 2A, 3A, 4A, 4C, 5A, 5C, 6A, 6C, 6F, 7A, 7C, 7E, 7G, 8A, 9A and 9C.

15

F.     Growth Assay

A growth assay, applicable to yeast strains Y190 and Y153b1 which carry both *GAL-HIS3* and *GAL-lacZ* reporters, was sometimes used as a complement to the color assay described above. Yeast transformants were streaked onto selective plates containing 40-50 mM  
20 3-AT and no Histidine, and incubated at 30°C for 3-7 days. Growth (corresponding to the level of *HIS3* expression) was monitored as an indicator of the interaction between fusion proteins. In cases where both assays were used, the amount of cell growth typically correlated well with the color intensity in the  $\beta$ -gal assay.

25           G.     Yeast Growth. Drug Treatment

Yeast were typically grown in YPD (rich non-selective) or synthetic complete (SC) medium with selected component drop-outs, depending on the plasmid introduced, following standard procedures (Sherman, *et al.*, Ausubel, *et al.*).

Experiments utilizing treatment with drugs or additives were performed by including  
30 the drug or additive in the medium. For plating, the agar was autoclaved, allowed to cool to 50°C, and the drug or additive was added before pouring the plates. Unless otherwise indicated, drugs and additives were added to result in the following final medium concentrations: FK506: 1  $\mu$ g/ml, cyclosporin A: 10  $\mu$ g/ml, rapamycin: 10 ng/ml, and hygromycin B: 40  $\mu$ g/ml.

## H. Antibodies

Polyclonal and monoclonal antibodies, for use in the present invention, can be prepared by standard methods (Harlow, *et al.*) utilizing the CNI polypeptides of the present invention, for example, a substantially purified CNI/ $\beta$ -galactosidase fusion protein (Example 9).  
5 Antibodies can also be generated by recombinant techniques (Cabilly, *et al.*; Better, *et al.*; Skerra, *et al.*). In addition to whole antibody molecules, antibody fragments retaining the immunological specificity of the whole antibody may also be used in the practice of the present invention (*e.g.*, Fab and F(ab')<sub>2</sub> fragments of IgG (Pierce Chemical, Rockford, IL)). The antibodies can be purified by standard methods to provide antibody preparations which are  
10 substantially free of serum proteins that may affect reactivity (*e.g.*, affinity purification (Harlow *et al.*)).

### EXAMPLE 1

#### Isolation of CN1c

##### 15 A. Library Screening

Yeast strain Y190 was transformed with pGBT-CNA1 *TRP1* (GBT-A1) hybrid plasmid using the transformation protocol described by Schiestl, *et al.* Transformants were selected, colony purified, and a single transformant was selected to make (Y190 GBT-A1)-competent cells, following the procedure described in Guthrie, *et al.*.

20 The three pGAD yeast fusion libraries described above, carrying fusions between G4AD and yeast genomic DNA *Sau3A*I fragments in each reading frame, were then used to transform (Schiestl, *et al.*) the Y190 GBT-A1-containing cells. Transformants were plated onto SC-Trp-Leu-His plates containing 40 mM 3-aminotriazole (3-AT; Sigma Chemical Co., St. Louis, MO) and incubated at 30°C for 6 days to screen for HIS<sup>+</sup> colonies (Durfee, *et al.*).

25 His<sup>+</sup> colonies were replica plated onto nitrocellulose filters (Schleicher & Schuell, Keene, NH), frozen in liquid nitrogen for approximately 30 seconds, and incubated at 30°C for 12 hours with Z buffer (see above) containing the chromogenic substrate X-Gal (0.1%) to assay  $\beta$ -gal activity (Breedon, *et al.*).

Candidate positive (blue) colonies were re-streaked for single colonies. Single colonies  
30 were purified and retested using the above protocol. Colonies which reproducibly tested positive were screened using PCR with primers directed against the internal portion of GAL-4 (*i.e.* the portion between the DNA binding domain and the activation domain). The sequences of the primers, G4-PCR-A and G4-PCR-B, are given as SEQ ID NO:22 and SEQ ID NO:23,

respectively. Colonies yielding a PCR product were identified as containing intact GAL4, and were eliminated.

The GBT-A1 *TRP*<sup>+</sup> plasmid was eliminated by growing in *Trp*<sup>+</sup> liquid media for 2-3 days, plating on -Leu media and then replica-plating on -Leu and -Trp plates to identify and eliminate colonies that had lost the GBT-A1 plasmid, yet still gave a positive signal.

Plasmid DNA was extracted from the remaining *Leu*<sup>+</sup> candidates. The plasmid DNA was transformed into *E coli* JBe181 and plated on -Leu media to select for library plasmids. The library plasmids isolated by this method were introduced back to the yeast reporter strains either alone or with test G4BD fusions: GBT, GBT-A1, and AS-lamin.

A parallel specificity assay was conducted by mating. Candidate strains, as described above, were 3-AT growth positive and X-gal positive when both the library and GBT-A1 plasmids were present. After elimination of the GBT-A1 plasmid from these strains, strains that were *Leu*<sup>+</sup> *Trp*<sup>-</sup> 3-AT growth<sup>-</sup> and  $\beta$ -gal<sup>-</sup> were mated to the following strains: Y187 (*MAT $\alpha$* ) carrying GBT, GBT-A1, or AS-lamin, and the diploids were assayed.

Among the 3-AT positive,  $\beta$ -gal positive candidates identified by the secondary screening method just described, one clone (III-21S, later termed GAD-CN1c) was specifically positive in conjunction with GBT-A1 in both the transformation assay and the mating assay.

#### B. Sequence of CN1c

Clone III-21S was sequenced as above. The sequence is presented herein as SEQ ID NO:1, and a schematic representation of the clone is shown in Figure 1A. The *Sau3AI* library insert encodes 306 amino acids followed by a stop codon in frame with the coding sequence of GAL4, but does not contain an in-frame ATG start site upstream of the stop codon. This is consistent with the fragment encoding a c-terminal portion of a complete gene product. Accordingly, the clone was termed CN1c, with the lowercase "c" representing "c-terminal".

The stippled portion between GAL-4AD and CN1c in Figure 1A represents the linker discussed in Materials and Methods, above. Also indicated in the Figure are the approximate locations of in-frame start (ATG) and stop codons, and *Sau3AI* restriction sites.

#### C. Isolation of a Full Length Clone

A <sup>32</sup>P-labeled CN1c probe was generated from the 1.22 kb CN1c insert of clone III-21S by polymerase chain reaction (PCR) using primers represented as SEQ ID NO:7 and SEQ ID NO:8. The probe was used to map the gene to the right arm of chromosome 11 by hybridization screening (Sambrook, *et al.*) a panel of  $\lambda$  clones (American Type Culture

Collection (ATCC), Rockville, MD) spanning the entire yeast genome. Two clones, 70500 and 70590, gave positive hybridization signals. A phage lysate of clone 70500 in  $\lambda$  MG3 was obtained from the ATCC, was amplified, purified, restriction-mapped and used as a DNA source for subsequent cloning experiments.

5       The phage DNA was digested with *SacI*, yielding a 7.1 kb fragment containing the entire CNI gene. This fragment was cloned into "BLUESCRIPT SK-" (Stratagene, La Jolla, CA) cut with *SacI*, yielding plasmid CNI7.1. Plasmid CNI7.1 was digested with *MunI* and *EcoRV*, releasing a 3.16 kb fragment containing the entire coding sequence of CNI. The 3.16 kb fragment was then cloned into each of YEp352(HIS), YEp352(TRP), YEp352(URA), and  
10 "BLUESCRIPT SK-", each cut with *EcoRI* and *SmaI*, yielding plasmids CNIH, CNIT, CNIU and CNI3.2, respectively. The sequence of the 3.16 kb *MunI/EcoRV* fragment is encompassed by the 3.5 kb sequence presented as SEQ ID NO:4. The *MunI* site is at nucleotide 100 of SEQ ID NO:4, and the *EcoRV* site is at nucleotide 3263 of SEQ ID NO:4.

A schematic diagram of the sequence presented as SEQ ID NO:4 is shown in Figure  
15 1B. This sequence contains the 3.16 kb *MunI/EcoRV* fragment used in many of the experiments described herein (depicted in Figure 1B as the portion between the *MunI* and *EcoRV* sites), which contains the entire 2.75 kb coding sequence of CNI (SEQ ID NO:6; graphically indicated in Figure 1B by the locations of the "ATG" and "Stop" codons).

Figure 13 also shows the location of certain features of the sequence. For example,  
20 "PEST" motifs (Rogers, *et al.*, Dice) are indicated by bars over the corresponding sequence.

A search of known DNA and protein sequences turned up no obvious matches or homologies to genes in other organisms. Accordingly, CNI may represent a new type of calcineurin-binding protein.

25

## EXAMPLE 2

### Binding of CNIC to Calcineurin

Y190 yeast carrying the plasmids indicated below were assayed for  $\beta$ -gal activity by color development assay described above to determine the specificity of binding of CNIC to subunits of calcineurin.

30       Exemplary data, in the form of images of filters having yeast colony replicas that had undergone the  $\beta$ -gal color development assay are shown in Figure 2A. The legend for Fig. 2A is shown in Figure 2B. Numbers in the legend refer to locations of yeast colonies expressing particular combinations of plasmid constructs. The constructs are as follows: 20:GBT-A1 and

GAD-B1, 21:GBT-A1 and GAD-CN1c, 22:GBT-A1 and GAD-CN1c, 23:GBT-A1 and GAD-CN1c 24:GBT-A2 and GAD-CN1c 25:AS-lamin and GAD-CN1c 26:GBT and GAD-CN1c.

Yeast colonies used in the assay were derived by several different methods. Those at location 22 were purified colonies from the original library screen, those at 21 were colonies  
5 transformed with mini-prep DNA of the isolated GAD-CN1c plasmid, and the remaining colonies (23, 24, 25 and 26) were transformed with maxi-prep (Qiagen, Chatsworth, CA) DNA of GAD-CN1c.

A comparison of the intensities of the blue  $\beta$ -gal reaction product indicates that CN1c interacted strongly with CNA1 regardless of the source of the CN1c plasmid DNA (20, 21, 22  
10 and 23), and somewhat less strongly with CNA2 (24). Neither cells containing AS-lamin with GAD-CN1c (25), nor cells containing only GBT with GAD-CN1c (26) showed a detectable signal above background. Two subunits of calcineurin (CNA1, CNB1) known to interact were used as a positive control for the assay (20).

In summary, the data above show that CN1c interacted specifically with CNA1 and  
15 CNA2, but not with G4BD or lamin C.

A similar set of experiments was conducted using constitutively-active CNA subunits, as well as calcineurin subunit CNB1. As described in Materials and Methods, above, CNA1 $\Delta$ C and CNA2 $\Delta$ C were each missing a C-terminal portion of the protein containing an autoinhibitory domain. Exemplary results from these experiments are shown in Figure 3A.  
20 The legend for Fig. 3A is shown in Figure 3B. Locations of yeast colonies expressing specific constructs are as follows: 27:GBT and GAD-CN1c, 28:GBT-A1 $\Delta$ C and GAD-CN1c, 29:GBT-A2 $\Delta$ C and GAD-CN1c, 30:GBT-A1 $\Delta$ C and GAD, 31:GBT-A2 $\Delta$ C and GAD, 32:GBT-B1 and GAD-CN1c, 33:GBT-B1 and GAD, 34:GBT-A1 $\Delta$ C and GAD-B1, and 35:GBT-A2 $\Delta$ C and GAD-B1.

25 The data show that GBT-A1 $\Delta$ C and GAD-CN1c (28) gave a definite positive signal, while GBT-A2 $\Delta$ C and GAD-CN1c (29) was weaker, though still detectable above its background (*i.e.* GBT-A2 $\Delta$ C and GAD; 31). The signal from GBT-B1 and GAD-CN1c (32) was not detectable above vector background (GBT-B1 and GAD; 33). Positive controls GBT-A1 $\Delta$ C and GAD-B1 (34) and GBT-A2 $\Delta$ C and GAD-B1 (35) gave strong signals.

30 These data show that CN1c interacted specifically with CNA1 $\Delta$ C and CNA2 $\Delta$ C, but not with CNB1.



**EXAMPLE 3****Effects of Immunosuppressant Drugs on Binding of CN1c to Calcineurin in B1<sup>™</sup>, B1  
Deletion and B1 Overproducing Yeast Strains**

Three yeast strains carrying the plasmids indicated below were assayed for  $\beta$ -gal activity as above to determine if the immunosuppressant drugs FK506, cyclosporin A (CsA) and rapamycin affect the binding of CN1c to subunits of calcineurin. The experiments were performed in yeast strains wild-type for the CNB1 subunit, null for the CNB1 subunit, and in yeast transformed with a high efficiency expression vector containing DNA encoding the CNB1 subunit.

Exemplary data, in the form of filter images produced as above are shown in Figures 4A, 4C, 5A and 5C. Plates used to make the filters shown in Figs. 4A and 4C were replicas from one master plate, while plates used to make the filters shown in Figs. 5A and 5C were replicas from another plate. The plates used to generate filters shown in Figs. 4A and 5A were without FK506, while the plates used to generate filters shown in Figs. 4C and 5C contained 1  $\mu$ g/ml FK506.

The legends for Figs. 4A and 4C are shown in Figures 4B and 4D, respectively. Since the imaged filters in Figs. 4A and 4C were from replica plates, corresponding locations on each filter contain material from the same yeast colonies. Accordingly, the locations are referred to by the same "base" numbers in the legends. To facilitate reference to a specific location on a specific filter, the base numbers are followed by a lowercase letter that is different for each of the individual filters. For example, in the present figure, "a" follows the base numbers to identify locations on the filter shown in Fig. 4A, while a "b" follows the base numbers to identify locations on the plate in Fig. 4C. This labeling scheme is used in other experiments detailed herein where multiple filter lifts are shown.

The interactions of various combinations of proteins expressed by constructs indicated below was studied in three yeast strains. Strain Y153b<sup>+</sup>, at 36-40, is null for the CNB1 subunit of calcineurin. Strains Y190 (41-46) and Y526 (47-51) are wild-type for CNA1, CNA2 and CNB1.

Hybrid proteins expressed by colonies at specific locations are as follows: 36:GBT-A1 and GAD-CN1c, 37:GBT-A2 and GAD-CN1c, 38:GBT-A1 $\Delta$ C and GAD-CN1c, 39:GBT and GAD-CN1c, 40:GBT-A2 $\Delta$ C and GAD-CN1c, 41:GBT-A1 and GAD-CN1c, 42:GBT and GAD-CN1c, 43:GBT-A2 and GAD-CN1c, 44:GBT-A1 $\Delta$ C and GAD-CN1c, 45:GBT-A1 and GAD-B1, 46:GBT-A2 $\Delta$ C and GAD-CN1c, 47:GBT-A1 and GAD-CN1c, 48:GBT and GAD-CN1c,

49:GBT-A2 and GAD-CNIc, 50:GBT-A1ΔC and GAD-CNIc, and 51:GBT-A2ΔC and GAD-CNIc.

Yeast strain Y526 was used for all experiments shown in Figs. 5A, 5B, 5C and 5D. The expression vector B1/YEp352(HIS) was not used in strains Y190 or Y153b1 because they are HIS<sup>+</sup> in the absence of 3-AT.

The base numbers in Figures 5B and 5D correspond to locations of colonies expressing the following constructs: 52:GBT-A1, GAD-CNIc and YEp352, 53:GBT-A1, GAD-CNIc and B1/YEp352, 54:GBT-A2, GAD-CNIc and YEp352, 55:GBT-A2, GAD-CNIc and B1/YEp352, 56:GBT, GAD-CNIc and B1/YEp352, 57:GBT-A1ΔC, GAD-CNIc and B1/YEp352, and 58:GBT-A2ΔC, GAD-CNIc and B1/YEp352.

A comparison of data shown in Fig. 4A (no added drugs) shows the effect of deleting the endogenous host CNB1 gene on interactions between CNIc and calcineurin subunits CNA1, CNA2, CNA1ΔC and CNA2ΔC. Note that interactions in panels 36a-40a (CNB1 null strain) were all stronger (with the exception of the negative control in 39) than interactions in corresponding panels 41a-51a (strains wild-type for CNB1). This result indicates that interaction between CNIc and CNA subunits were enhanced by the deletion of the CNB1 subunit.

Comparison of corresponding panels in Figs. 4A and 4C shows the effects of FK506 on CNIc-CNA/CNAΔC interactions. The drug enhanced interactions under all except control (39, 42 and 48) conditions. The effect was most striking in yeast strains wild-type for the CNB1 subunit (*e.g.*, compare 50a with 50b, and 51a with 51b).

The drug also markedly enhanced CNIc-CNA/CNAΔC interactions under conditions where the CNB1 subunit was overexpressed. Figure 5A shows the effect of overexpressing CNB1 on CNIc-CNA/CNAΔC interactions in the absence of drug. Colonies expressing B1/YEp352 (53a, 55a-58a) had reduced signal as compared with controls expressing only YEp352 (52a, 54a). Inclusion of FK506 in the plating medium (Fig. 5C), however, enhanced interactions in all colonies, except the negative controls (56).

Taken together, the above data demonstrate that the interaction of CNIc with CNA and CNAΔC was markedly enhanced by FK506. The interaction was also enhanced by deletion of CNB1 and diminished by overexpression of CNB1, and the inhibitory effect of CNB1 overproduction was overcome by the stimulatory effect of FK506.

In light of the effects of FK506 on CNIc-CNA/CNAΔC interactions, two other immunosuppressants, cyclosporin A and rapamycin, were examined in similar experiments. The results of these experiments are shown in Figures 6A-6F and 7A-7H. Filters shown in

Figs. 6A-6F were from replica plates, as were those in Figs. 7A-7H. Colonies shown in Figs. 6A and 7A were plated without drugs; those in Figs. 6C and 7C were plated with FK506 (1  $\mu$ g/ml), those in Figs. 6E and 7E with CsA (10  $\mu$ g/ml), and those in Fig. 7G with rapamycin (10 ng/ml). Yeast strains used were as follows: In Figs. 6A-6F, panels 59-63 were Y153b1, 64-69 were Y190, and 70-74 were Y526. In Figs. 7A-7H, panels 77-83 were Y526, and panels 75 and 76 were Y190.

The base numbers in Figures 6B, 6D and 6F correspond to locations of colonies expressing the following constructs: 59:GBT-A1 and GAD-CN1c, 60:GBT-A2 and GAD-CN1c, 61:GBT-A1 $\Delta$ C and GAD-CN1c, 62:GBT and GAD-CN1c, 63:GBT-A2 $\Delta$ C and GAD-CN1c, 64:GBT-A1 and GAD-CN1c, 65:GBT and GAD-CN1c, 66:GBT-A2 and GAD-CN1c, 67:GBT-A1 $\Delta$ C and GAD-CN1c, 68:GBT-A1 and GAD-B1, 69:GBT-A2 $\Delta$ C and GAD-CN1c, 70:GBT-A1 and GAD-CN1c, 71:GBT and GAD-CN1c, 72:GBT-A2 and GAD-CN1c, 73:GBT-A1 $\Delta$ C and GAD-CN1c, and 74:GBT-A2 $\Delta$ C and GAD-CN1c.

The base numbers in Figures 7B, 7D, 7F and 7H correspond to locations of colonies expressing the following constructs: 75:GBT-FKBP and GAD, 76:GBT-FKBP and GAD-CN1c, 77:GBT-A1, GAD-CN1c and YEp352, 78:GBT-A1, GAD-CN1c and B1/YEp352, 79:GBT-A2, GAD-CN1c and YEp352, 80:GBT-A2, GAD-CN1c and B1/YEp352, 81:GBT, GAD-CN1c and B1/YEp352, 82:GBT-A1 $\Delta$ C, GAD-CN1c and B1/YEp352, and 83:GBT-A2 $\Delta$ C, GAD-CN1c and B1/YEp352.

The data presented in Figures 6A and 6C are essentially equivalent to those presented in Figures 4A and 4C, respectively. The constructs and yeast strains at corresponding locations were the same. As expected, the  $\beta$ -gal signal was also essentially equivalent between the two sets. Data shown in Figure 6E demonstrate that like FK506, the immunosuppressant cyclosporin A was also effective in enhancing interaction of CN1c with CNA and CNA $\Delta$ C. Both FK506 and cyclosporin A are known to exert their immunosuppressive effects through inhibition of calcineurin activity (Cyert).

Similarly, data shown in Figures 7A and 7C are essentially equivalent to those in Figures 5A and 5C, except that a top panel has been added in Figs. 7A-H. As above, the corresponding panels show the same constructs and yeast strains. The added panels (75 and 76) assessed the interaction of an FK506 binding protein (FKBP) with CN1c, and indicate that there were no detectable interactions between these proteins. Results in Fig. 7E demonstrate that cyclosporin A had a similar effect to FK506 in cells overexpressing CNB1 -- that is, it enhanced the interactions between CN1c and CNA/CNA $\Delta$ C.

In contrast, data presented in Fig. 7G show that the immunosuppressant rapamycin, which is known not to target calcineurin, had no detectable effect on CN1c-CNA/CNAΔC interactions (compare Fig. 7G with Fig. 7A).

Taken together, the above data show that like FK506, cyclosporin A (CsA), but not rapamycin, also enhanced the interaction of CN1c with CNA and CNAΔC. CN1c didn't interact with FKBP with or without FK506.

#### EXAMPLE 4

##### Effects of CNI on FKBP/FK506 binding to Calcineurin

Y526 cells, carrying the plasmids indicated below, were grown in -Trp-Leu-His liquid media with or without FK506 (1 μg/ml) until OD<sub>600</sub> reached about 1.0. Approximately the same number of cells, calculated based on OD<sub>600</sub> and equivalent to 1 ml of an OD<sub>600</sub>=1 suspension, was harvested from each culture, washed once with ddH<sub>2</sub>O, centrifuged briefly, and the pellet was resuspended in 30 μl ddH<sub>2</sub>O and transferred onto a nitrocellulose filter. The filters were frozen in liquid nitrogen as described above, placed in a 8.5 cm petri dish containing a sheet of Whatman No. 1 paper (Whatman International LTD, Maidstone, UK) in 1.6 ml Z buffer containing 0.1% X-Gal, and incubated at 30°C for 8 hours.

Figure 8A presents exemplary data from studies to assess the effect of CNI overexpression on FK506-mediated FKBP interactions with CNA2. The legend for Fig. 8A is shown in Figure 8B. Locations of yeast colonies expressing specific constructs: 84:GBT-FKBP, GAD-A2 and YEp352, 85:GBT-FKBP, GAD-A2 and YEp352, 86:GBT-FKBP, GAD-A2 and CNI/YEp352, and 87:GBT-FKBP, GAD-A2 and CNI/YEp352. The cells at 85 and 87 were exposed to FK506, while those at 84 and 86 were not.

Data in Figure 8A demonstrate that, in the absence of FK506, FKBP and CNA2 showed no detectable interaction (84). In the presence of FK506, however, the proteins interacted (85), presumably because FK506 formed a complex with FKBP, which then bound CNA2 (Cyert).

Data in Figure 8A further show that, in the absence of FK506, CNI had no effect on the lack of interaction between FKBP and CNA2 (86). In the presence of FK506 (87), however, CNI potentiated, or enhanced the binding between FKBP and CNA2 (compare the intensity at 87 with that at 85). This effect suggests that CNI and similar compounds may be employed to increase the sensitivity of calcineurin to immunosuppressant drugs that act on it, and in this way, decrease the amount of the immunosuppressant required for a particular level of immunosuppression.

Experiments illustrated in Figures 9A and 9B demonstrate that CNI overproduction had little or no effect on the binding of CNA2 to CNB1, providing support for the specificity of the stimulatory effect that CNI overproduction had on the FK506-dependent binding of FKBP to calcineurin. The legends for Fig. 9A and 9B are shown in Figures 9B and 9D, respectively.

- 5 Locations of yeast colonies expressing the following constructs: 88:GBT-A2, GAD-B1 and YEp352, 89:GBT-A2, GAD-B1 and YEp352, 90:GBT-A2, GAD-B1 and CNI/YEp352, and 91:GBT-A2, GAD-B1 and CNI/YEp352. The colonies at 89 and 91 were exposed to FK506, while colonies at 88 and 90 were not.

- 10 Taken together, the above data show that overexpression of the full-length CNI clone markedly enhanced the FK506-dependent interaction of FKBP with CNA, although it didn't affect the interaction between CNA and CNB1.

#### EXAMPLE 5

##### Co-Immunoprecipitation of CNIC and CNA

- 15 Yeast BJ2407 harboring AS-CNIC, which carries an influenza hemagglutinin (HA) epitope tag (Wilson, *et al.*), and GAD-A2 (lanes 1, 3) or A2/YEp352 (lanes 2, 5), and strain MCY300-1 (*cna1<sup>-</sup>cna2<sup>-</sup>*; lane 4) were grown in selective media to  $OD_{600}=0.8$ . The cells were harvested, lysed, and immunoprecipitated in the presence of 25  $\mu\text{g/ml}$  FK506 with anti-HA monoclonal antibody (obtained from M. Kirschner, Harvard University, Boston, MA; Wilson, *et al.*), following protocols described in Harlow, *et al.* The cell extracts (lanes 3-5) and the  
20 immune complex (lanes 1, 2) were resolved by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE; Laemmli) followed by western blot with a rabbit anti-CNA2 polyclonal antibody generated using standard methods (Harlow, *et al.*). Bound anti-CNA2 antibody was visualized with the "ECL" kit (Amersham, Arlington Heights, IL) using goat anti-  
25 rabbit antibody. Molecular weight markers are indicated on the right in kD.

The results demonstrate that CNIC was capable of binding to CNA2 tightly enough for the complex to be co-immunoprecipitated. This independent, biochemical assay confirmed the results described above obtained using the two hybrid protein interaction assay — that is, that CNIC physically interacted with and bound CNA subunits.

- 30 Cell extracts of BJ2407 harboring AS-CNIC, and GAD-A2 or A1/Yep351, and Y153b1 harboring AS-CNIC and GAD-A1 were subjected to SDS-PAGE followed by western blot with anti HA antibody. The results showed that CNIC was present at very low levels *in vivo*.

The results are consistent with the observations that a limited amount of CNA2 was precipitated by anti-HA antibody recognizing the CN1c fusions, and that CN1 contains PEST-like motifs, a feature of proteins with a short half-life *in vivo* (Rogers, *et al.*).

5

#### EXAMPLE 6

##### Northern Blot of CN1c

Norther blots (*e.g.*, Sambrook, *et al.*) of yeast total RNA were hybridized with a CN1c probe. Exemplary data are shown in Figure 11. 20 µg yeast RNA from YPH499 (lane 1) and MCY300-1 (lane 2) was resolved in a formaldehyde-agarose gel, transferred onto "HYBOND  
10 N+" membrane (Amersham, Arlington Heights, IL), and hybridized with 5x10<sup>6</sup> cpm/ml probe of the CN1c insert (1.22kb). A single message of approximately 2.9 kb was detected in both strains at about the same level following an 18-hour exposure on XAR5 film (Eastman Kodak, Rochester, NY).

The data indicate that CN1 was a physiologically expressed gene encoding a 2.9 kb  
15 message in yeast.

#### EXAMPLE 7

##### Chromosome Mapping of CN1c

A yeast chromosome blot obtained from the ATCC was hybridized with probe of the  
20 CN1c insert following the Southern hybridization procedure described in Sambrook, *et al.* A positive hybridization signal was obtained with two ATCC yeast genomic λ clones derived from chromosome 11. Clone 70500 had a relatively strong signal, while clone 70590 had a somewhat weaker one. A phage lysates of clone 70500 was ordered from the ATCC, amplified, purified, restriction-mapped, and used as a DNA source for cloning full length CN1  
25 (Example 1).

#### EXAMPLE 8

##### CN1 null Mutants

###### 1. Construction of *cni* Null Mutation

30 A 5', 1.8 kb *Bgl*II-*Hind*III and a 3', 0.9 kb *Xba*I-*Bgl*II fragment of CN1 were ligated into pRS305(LEU2) (Sikorski, *et al.*). The resultant plasmid had a deletion of a 2 kb *Hind*III-*Xba*I fragment from the coding sequence of CN1. This *cni::LEU2* mutant was introduced into the genomes of yeast haploid strains YPH499 (Sikorski, *et al.*), MCY300-1 (*cna1<sup>-</sup> cna2<sup>-</sup>*) and DD12 (*cnb1<sup>-</sup>*) (Cyert, *et al.*, 1991, Cyert, *et al.*, 1992) as well as two diploid strains.

Leucine prototrophs were isolated at high frequency from all strains, and hybridization analysis confirmed that the *cni::LEU2* allele had replaced the CNI gene. The experiments indicate that CNI is not essential for viability, since CNI deletion mutant strains (even *cni<sup>-</sup> cni<sup>-</sup>* double mutants) can survive.

5        *CNI* was deleted from three yeast strains: YPH499 (WT), MCY300-1 (*cna1<sup>-</sup>cna2<sup>-</sup>*), and DD12 (*cnb1<sup>-</sup>*), resulting in *cni<sup>-</sup>* strains LHy499, LHy300 and LHy12, respectively. Cells representing four colonies of each *cni* knockout strain and two colonies of each parent strain were grown in liquid YPD (Sherman, *et al.*) to saturation. Same numbers of cells from each culture were then plated onto YPD+Hygromycin B (40  $\mu$ g/ml) and growth was monitored at  
10    30°C.

CNI deletions in each strain rendered that strain more resistant to hygromycin B. The effect was particularly pronounced in both MCY300-1 and DD12, suggesting that CNI functions as a suppressor of CN mutant's sensitivity to hygromycin B. The data indicate that deletion of *CNI* results in higher resistance to hygromycin B.

15

#### EXAMPLE 9

##### Isolation of CNI/ $\beta$ -Galactosidase Fusion Protein

A CNI coding sequence is cloned into the  $\lambda$  gt11 vector (Stratagene, La Jolla, CA). The coding frame is cloned in-frame to the  $\beta$ -galactosidase coding sequences present in  $\lambda$  gt11.  
20    Bacterial lysogens infected either with lambda phage gt11 or with gt11/CNI are incubated in 32°C until the culture reaches to an O.D. of 0.4. Then the culture is incubated in a 43°C water bath for 15 minutes to induce gt11 peptide synthesis, and further incubated at 37°C for 1 hour. Bacterial cells are pelleted and lysed in lysis buffer (10 mM Tris, pH 7.4, 2 % "TRITON X-100" and 1% aprotinin). Bacterial lysates are clarified by centrifugation (10K,  
25    for 10 minutes, Sorvall JA20 rotor) and the clarified lysates are incubated with Sepharose 4B beads conjugated with anti- $\beta$ -galactosidase (Promega).

Binding and elution of  $\beta$ -galactosidase fusion proteins are performed according to the manufacturer's instruction. Typically binding of the proteins and washing of the column are done with lysis buffer. Bound proteins are eluted with 0.1 M carbonate/bicarbonate buffer,  
30    pH 10.

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: The Board of Trustees of the Leland Stanford Junior University
- (ii) TITLE OF INVENTION: Calcineurin Interacting Protein Compositions and Methods
- (iii) NUMBER OF SEQUENCES: 23
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Dehlinger & Associates
  - (B) STREET: 350 Cambridge Avenue, Suite 250
  - (C) CITY: Palo Alto
  - (D) STATE: CA
  - (E) COUNTRY: USA
  - (F) ZIP: 94306
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE: 23-OCT-1995
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 08/328,322
  - (B) FILING DATE: 24-OCT-1994
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Sholtz, Charles K.
  - (B) REGISTRATION NUMBER: P38,615
  - (C) REFERENCE/DOCKET NUMBER: 8600-0151.41
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (415) 324-0880
  - (B) TELEFAX: (415) 324-0960

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1222 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: Sau3AI fragment containing CN1c coding sequence
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..918



## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAT CAA AGT AGC AAT GTC TTC GCA TCC AAA CAG CTG GTC GCA AAC ATT	48
Asp Gln Ser Ser Asn Val Phe Ala Ser Lys Gln Leu Val Ala Asn Ile	
1 5 10 15	
TAT AAG CCC AAT CAG ATT CCA AGA GAA TTA ACT TCT CCT CAG GCG TTA	96
Tyr Lys Pro Asn Gln Ile Pro Arg Glu Leu Thr Ser Pro Gln Ala Leu	
20 25 30	
CCA TTA TCG CCC ATC ACC TCA CCA ATT CTC AAT TAC CAA CCA TTA TCA	144
Pro Leu Ser Pro Ile Thr Ser Pro Ile Leu Asn Tyr Gln Pro Leu Ser	
35 40 45	
AAC TCC CCG CCT CCA GAT TTT GAT TTT GAT CTA GCT AAG CGC GGC GCA	192
Asn Ser Pro Pro Pro Asp Phe Asp Phe Asp Leu Ala Lys Arg Gly Ala	
50 55 60	
GCC GAT TCT CAT GCT ATT CCT GTG GAT CCT CCA TCA TAT TTT GAT GTA	240
Ala Asp Ser His Ala Ile Pro Val Asp Pro Pro Ser Tyr Phe Asp Val	
65 70 75 80	
TTA AAG GCC GAT GGG ATT GAA TTG CCA TAC TAC GAT ACA AGT TCA TCT	288
Leu Lys Ala Asp Gly Ile Glu Leu Pro Tyr Tyr Asp Thr Ser Ser Ser	
85 90 95	
AAA ATT CCT GAA CTA AAA CTA AAC AAA TCT AGA GAG ACA TTG GCC AGC	336
Lys Ile Pro Glu Leu Lys Leu Asn Lys Ser Arg Glu Thr Leu Ala Ser	
100 105 110	
ATT GAG GAG GAC TCA TTC AAT GGT TGG TCT CAA ATT GAT GAC TTA TCC	384
Ile Glu Glu Asp Ser Phe Asn Gly Trp Ser Gln Ile Asp Asp Leu Ser	
115 120 125	
GAC GAA GAT GAC AAT GAT GGC GAT ATA GCA TCT GGT TTC AAC TTC AAG	432
Asp Glu Asp Asp Asn Asp Gly Asp Ile Ala Ser Gly Phe Asn Phe Lys	
130 135 140	
CTG TCA ACC AGT GCT CCG AGT GAG AAC GTT AAT TCA CAC ACT CCT ATT	480
Leu Ser Thr Ser Ala Pro Ser Glu Asn Val Asn Ser His Thr Pro Ile	
145 150 155 160	
TTG CAG TCT TTA AAC ATG AGT CTT GAT GGG AGA AAA AAA AAT CGT GCC	528
Leu Gln Ser Leu Asn Met Ser Leu Asp Gly Arg Lys Lys Asn Arg Ala	
165 170 175	
AGT CTA CAC GCA ACA TCA GTG TTA CCT AGT ACA ATA AGA CAG AAC AAT	576
Ser Leu His Ala Thr Ser Val Leu Pro Ser Thr Ile Arg Gln Asn Asn	
180 185 190	
CAG CAT TTC AAT GAC ATA AAC CAG ATG CTA GGC AGT AGT GAC GAA GAT	624
Gln His Phe Asn Asp Ile Asn Gln Met Leu Gly Ser Ser Asp Glu Asp	
195 200 205	
GCC TTT CCC AAA AGC CAA TCA TTA AAT TTC AAT AAG AAA CTA CCA ATA	672
Ala Phe Pro Lys Ser Gln Ser Leu Asn Phe Asn Lys Lys Leu Pro Ile	
210 215 220	
CTT AAA ATT AAT GAT AAC GTC ATA CAA TCA AAC AGC AAT AGT AAT AAC	720
Leu Lys Ile Asn Asp Asn Val Ile Gln Ser Asn Ser Asn Ser Asn Asn	
225 230 235 240	
AGA GTT GAT AAT CCA GAA GAT ACA GTG GAT TCT TCA GTC GAT ATT ACA	768
Arg Val Asp Asn Pro Glu Asp Thr Val Asp Ser Ser Val Asp Ile Thr	
245 250 255	

40

GCA TTT TAT GAT CCA AGA ATG TCA TCA GAT TCC AAA TTT GAT TGG GAG 816  
 Ala Phe Tyr Asp Pro Arg Met Ser Ser Asp Ser Lys Phe Asp Trp Glu  
                   260                                  265                                  270

GTA AGC AAG AAC CAT GTT GAC CCA GCA GCC TAC TCG GTT AAC GTT GCT 864  
 Val Ser Lys Asn His Val Asp Pro Ala Ala Tyr Ser Val Asn Val Ala  
                   275                                  280                                  285

AGT GAA AAC CGT GTA CTG GAC GAC TTT AAG AAA GCA TTT CGC GAA AAG 912  
 Ser Glu Asn Arg Val Leu Asp Asp Phe Lys Lys Ala Phe Arg Glu Lys  
                   290                                  295                                  300

AGA AAA TAAGTACATT ATTTTCATTC TCCGACAGAA TTGCTACCAT TTTACTTTGT 968  
 Arg Lys  
 305

GTCTGTGAT TCAATAGTGT ACAATATATT GGACATTTTA TAGTATACAA ATATACACCA 1028  
 TCAATCTATA CATCCATATC ACTTGTCGTA AAGATATCCC TTTTAAATAG TACAGCGATT 1088  
 AAAAAAATAA CATGATTAAC GTTCAGTTAC CAATGAGCTT ATTTATTAGG CTTGCTTTAG 1148  
 ATTTTTCCAA GTCAATTTTT GTTTTTTCTA ACGCTTGCAA CCTCATCTCA ACCTTCTTCC 1208  
 TTTGCAAGCA GATC 1222

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 306 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Asp Gln Ser Ser Asn Val Phe Ala Ser Lys Gln Leu Val Ala Asn Ile  
 1                  5                                  10                                  15

Tyr Lys Pro Asn Gln Ile Pro Arg Glu Leu Thr Ser Pro Gln Ala Leu  
                   20                                  25                                  30

Pro Leu Ser Pro Ile Thr Ser Pro Ile Leu Asn Tyr Gln Pro Leu Ser  
                   35                                  40                                  45

Asn Ser Pro Pro Pro Asp Phe Asp Phe Asp Leu Ala Lys Arg Gly Ala  
                   50                                  55                                  60

Ala Asp Ser His Ala Ile Pro Val Asp Pro Pro Ser Tyr Phe Asp Val  
                   65                                  70                                  75                                  80

Leu Lys Ala Asp Gly Ile Glu Leu Pro Tyr Tyr Asp Thr Ser Ser Ser  
                   85                                  90                                  95

Lys Ile Pro Glu Leu Lys Leu Asn Lys Ser Arg Glu Thr Leu Ala Ser  
                   100                                  105                                  110

Ile Glu Glu Asp Ser Phe Asn Gly Trp Ser Gln Ile Asp Asp Leu Ser  
                   115                                  120                                  125

Asp Glu Asp Asp Asn Asp Gly Asp Ile Ala Ser Gly Phe Asn Phe Lys  
                   130                                  135                                  140

Leu Ser Thr Ser Ala Pro Ser Glu Asn Val Asn Ser His Thr Pro Ile  
                   145                                  150                                  155                                  160

41

Leu Gln Ser Leu Asn Met Ser Leu Asp Gly Arg Lys Lys Asn Arg Ala  
                             165                            170                            175  
 Ser Leu His Ala Thr Ser Val Leu Pro Ser Thr Ile Arg Gln Asn Asn  
                             180                            185                            190  
 Gln His Phe Asn Asp Ile Asn Gln Met Leu Gly Ser Ser Asp Glu Asp  
                             195                            200                            205  
 Ala Phe Pro Lys Ser Gln Ser Leu Asn Phe Asn Lys Lys Leu Pro Ile  
                             210                            215                            220  
 Leu Lys Ile Asn Asp Asn Val Ile Gln Ser Asn Ser Asn Ser Asn Asn  
                             225                            230                            235                            240  
 Arg Val Asp Asn Pro Glu Asp Thr Val Asp Ser Ser Val Asp Ile Thr  
                             245                            250                            255  
 Ala Phe Tyr Asp Pro Arg Met Ser Ser Asp Ser Lys Phe Asp Trp Glu  
                             260                            265                            270  
 Val Ser Lys Asn His Val Asp Pro Ala Ala Tyr Ser Val Asn Val Ala  
                             275                            280                            285  
 Ser Glu Asn Arg Val Leu Asp Asp Phe Lys Lys Ala Phe Arg Glu Lys  
                             290                            295                            300  
 Arg Lys  
 305

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 918 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: CINC coding sequence

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GATCAAAGTA GCAATGTCTT CGCATCCAAA CAGCTGGTGC CAAACATTTA TAAGCCCAAT	60
CAGATTCCAA GAGAATTAAC TTCTCCTCAG GCGTTACCAT TATCGCCCAT CACCTCACCA	120
ATTCTCAATT ACCAACCATT ATCAAATCC CCGCCTCCAG ATTTTGATTT TGATCTAGCT	180
AAGCGCGGCG CAGCCGATTC TCATGCTATT CCTGTGGATC CTCCATCATA TTTTGATGTA	240
TTAAAGGCCG ATGGGATTGA ATTGCCATAC TACGATACAA GTTCATCTAA AATTCCTGAA	300
CTAAACTAA ACAATCTAG AGAGACATTG GCCAGCATTG AGGAGGACTC ATTCAATGGT	360
TGGTCTCAA TTGATGACTT ATCCGACGAA GATGACAATG ATGGCGATAT AGCATCTGGT	420
TTCAACTTCA AGCTGTCAAC CAGTGCTCCG AGTGAGAACG TTAATTCACA CACTCCTATT	480
TTGCAGTCTT TAAACATGAG TCTTGATGGG AGAAAAAAA ATCGTGCCAG TCTACACGCA	540

ACATCAGTGT	TACCTAGTAC	AATAAGACAG	AACAATCAGC	ATTTCAATGA	CATAAACCAG	600
ATGCTAGGCA	GTAGTGACGA	AGATGCCTTT	CCCAAAAGCC	AATCATTA	TTTCAATAAG	660
AAACTACCAA	TACTTAAAAT	TAATGATAAC	GTCATACAAT	CAAACAGCAA	TAGTAATAAC	720
AGAGTTGATA	ATCCAGAAGA	TACAGTGGAT	TCTTCAGTCG	ATATTACAGC	ATTTTATGAT	780
CCAAGAATGT	CATCAGATTC	CAAATTTGAT	TGGGAGGTAA	GCAAGAACCA	TGTTGACCCA	840
GCAGCCTACT	CGGTTAACGT	TGCTAGTGAA	AACCGTGTAC	TGGACGACTT	TAAGAAAGCA	900
TTTCGCGAAA	AGAGAAAA					918

(2) INFORMATION FOR SEQ ID NO:4:

**(i) SEQUENCE CHARACTERISTICS:**

- (A) LENGTH: 3500 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

(iii) **HYPOTHETICAL: NO**

(iv) ANTI-SENSE: NO

(v1) ORIGINAL SOURCE:

- (C) INDIVIDUAL ISOLATE: genomic DNA fragment containing full CNI coding sequence

**(ix) FEATURE:**

- (A) NAME/KEY: CDS  
(B) LOCATION: 376..3120

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CCGAACACTT	CCTTCGAGAG	AGTGCATTTT	ACTATGTGAA	CCAATTTTTC	CTCTTTTTCG	60										
GTTTGCAAGT	TCACCTGAAA	AACTGCTTAA	CACTACTAGC	AATTGCCCTA	TTGTCGTACG	120										
AGGACTTTGC	CAAATGTATT	CCCGGCTGTT	TGTAGTATAT	ATACGCAGAT	ATATAATAGC	180										
GCCGTCTTTT	TACCTCTTTG	AGCGAATTGC	CAAATATTGA	CTCTTTTGTC	TTATTTTCGCT	240										
ATCCCCATCT	TATCAAAAAT	GGGAACAACT	CGTTGAAATA	AGAGACAAGC	AACAAGAAAG	300										
ACAACCAACA	GAAAGTTCCA	TTCCGCACAA	ATACGCTGGA	ATCCCATAGA	ATATTGCTTG	360										
TTCCTCTATG	ACTAC	ATG	CTC	CAA	TTC	AAT	ACA	GAA	AAT	GAT	ACT	GTA	GCT	411		
	Met	Leu	Gln	Phe	Asn	Thr	Glu	Asn	Asp	Thr	Val	Ala				
	1				5						10					
CCA	GTG	TTT	CCC	ATG	GAG	CAA	GAT	ATA	AAT	GCA	GCA	CCT	GAT	GCC	GTC	459
Pro	Val	Phe	Pro	Met	Glu	Gln	Asp	Ile	Asn	Ala	Ala	Pro	Asp	Ala	Val	
		15					20					25				
CCA	CTG	GTG	CAG	ACA	ACA	ACA	CTA	CAA	GTC	TTT	GTA	AAG	CTT	GCC	GAA	507
Pro	Leu	Val	Gln	Thr	Thr	Thr	Leu	Gln	Val	Phe	Val	Lys	Leu	Ala	Glu	
	30					35					40					
CCC	ATA	GTG	TTT	TTA	AAA	GGA	TTT	GAA	ACT	AAC	GGA	CTG	TCT	GAA	ATA	555
Pro	Ile	Val	Phe	Leu	Lys	Gly	Phe	Glu	Thr	Asn	Gly	Leu	Ser	Glu	Ile	
	45				50					55					60	

GCC CCC AGT ATC TTA CGA GGA TCT CTT ATC GTC AGG GTG TTG AAA CCG Ala Pro Ser Ile Leu Arg Gly Ser Leu Ile Val Arg Val Leu Lys Pro 65 70 75	603
AAT AAA TTA AAA AGT ATA TCG ATA ACC TTC AAA GGA ATA TCC AGA ACA Asn Lys Leu Lys Ser Ile Ser Ile Thr Phe Lys Gly Ile Ser Arg Thr 80 85 90	651
GAG TGG CCG GAA GGT ATA CCA CCG AAG AGA GAA GAA TTT TCA GAT GTT Glu Trp Pro Glu Gly Ile Pro Pro Lys Arg Glu Glu Phe Ser Asp Val 95 100 105	699
GAA ACT GTT GTC AAT CAC ACA TGG CCA TTT TAT CAG GCG GAT GAC GGC Glu Thr Val Val Asn His Thr Trp Pro Phe Tyr Gln Ala Asp Asp Gly 110 115 120	747
ATG AAT TCT TTC ACC TTA GAA CAT CAC AGC TCA AAT AAT TCG TCC AAT Met Asn Ser Phe Thr Leu Glu His His Ser Ser Asn Asn Ser Ser Asn 125 130 135 140	795
CGC CCA TCT ATG AGC GAT GAA GAT TAT CTA CTT GAA AAA AGC GGT GCT Arg Pro Ser Met Ser Asp Glu Asp Tyr Leu Leu Glu Lys Ser Gly Ala 145 150 155	843
TCA GTA TAT ATC CCA CCA ACC GCT GAA CCC CCT AAA GAT AAT AGC AAT Ser Val Tyr Ile Pro Pro Thr Ala Glu Pro Pro Lys Asp Asn Ser Asn 160 165 170	891
CTA AGT CTG GAT GCC TAT GAG CGC AAC TCA TTG TCA TCC GAT AAT TTG Leu Ser Leu Asp Ala Tyr Glu Arg Asn Ser Leu Ser Ser Asp Asn Leu 175 180 185	939
AGT AAC AAG CCA GTA TCA AGT GAT GTT TCC CAT GAC GAC AGT AAA CTG Ser Asn Lys Pro Val Ser Ser Asp Val Ser His Asp Asp Ser Lys Leu 190 195 200	987
TTG GCT ATT CAA AAG ACA CCA TTA CCA TCA TCT AGT CGA AGA GGA TCG Leu Ala Ile Gln Lys Thr Pro Leu Pro Ser Ser Ser Arg Arg Gly Ser 205 210 215 220	1035
GTA CCG GCA AAT TTT CAC GGT AAC TCT TTG TCA CCT CAT ACC TTC ATA Val Pro Ala Asn Phe His Gly Asn Ser Leu Ser Pro His Thr Phe Ile 225 230 235	1083
TCT GAT TTG TTC ACA AAA ACA TTC AGT AAT AGT GGC GCT ACT CCA AGT Ser Asp Leu Phe Thr Lys Thr Phe Ser Asn Ser Gly Ala Thr Pro Ser 240 245 250	1131
CCT GAG CAA GAG GAT AAC TAT CTT ACA CCA TCC AAA GAT TCT AAA GAA Pro Glu Gln Glu Asp Asn Tyr Leu Thr Pro Ser Lys Asp Ser Lys Glu 255 260 265	1179
GTT TTT ATT TTT CGA CCG GGC GAT TAT ATT TAC ACT TTT GAA CAG CCA Val Phe Ile Phe Arg Pro Gly Asp Tyr Ile Tyr Thr Phe Glu Gln Pro 270 275 280	1227
ATA TCG CAA TCT TAT CCA GAA AGT ATA AAA GCC AAT TTT GGT TCC GTG Ile Ser Gln Ser Tyr Pro Glu Ser Ile Lys Ala Asn Phe Gly Ser Val 285 290 295 300	1275
GAG TAT AAA CTG TCA ATA GAC ATA GAG AGG TTT GGC GCA TTC AAA TCA Glu Tyr Lys Leu Ser Ile Asp Ile Glu Arg Phe Gly Ala Phe Lys Ser 305 310 315	1323
ACT ATA CAT ACT CAA TTA CCC ATC AAA GTC GTA AGG CTT CCT TCT GAT Thr Ile His Thr Gln Leu Pro Ile Lys Val Val Arg Leu Pro Ser Asp 320 325 330	1371

GGA TCC GTA GAA GAG ACT GAA GCT ATT GCA ATT TCC AAG GAC TGG AAA Gly Ser Val Glu Glu Thr Glu Ala Ile Ala Ile Ser Lys Asp Trp Lys 335 340 345	1419
GAT CTT CTT CAT TAT GAC GTG GTA ATT TTC TCG AAA GAG ATC GTT TTG Asp Leu Leu His Tyr Asp Val Val Ile Phe Ser Lys Glu Ile Val Leu 350 355 360	1467
AAT GCA TTT TTA CCC ATC GAT TTC CAT TTC GCT CCT CTA GAT AAA GTT Asn Ala Phe Leu Pro Ile Asp Phe His Phe Ala Pro Leu Asp Lys Val 365 370 375 380	1515
ACT CTG CAT CGT ATT AGA ATT TAT CTA ACA GAG TCT ATG GAA TAC ACT Thr Leu His Arg Ile Arg Ile Tyr Leu Thr Glu Ser Met Glu Tyr Thr 385 390 395	1563
TGT AAT AGT AAT GGA AAT CAC GAG AAG GCT CGT AGA TTA GAG CCA ACT Cys Asn Ser Asn Gly Asn His Glu Lys Ala Arg Arg Leu Glu Pro Thr 400 405 410	1611
AAA AAG TTT CTG TTG GCT GAA CAT AAC GGT CCT AAA CTG CCT CAT ATA Lys Lys Phe Leu Leu Ala Glu His Asn Gly Pro Lys Leu Pro His Ile 415 420 425	1659
CCA GCT GGT TCG AAT CCT TTG AAG GCT AAA AAT AGA GGG AAC ATC CTC Pro Ala Gly Ser Asn Pro Leu Lys Ala Lys Asn Arg Gly Asn Ile Leu 430 435 440	1707
TTG GAT GAA AAA TCC GGC GAT CTA GTT AAC AAA GAT TTT CAG TTC GAG Leu Asp Glu Lys Ser Gly Asp Leu Val Asn Lys Asp Phe Gln Phe Glu 445 450 455 460	1755
GTG TTT GTC CCA AGC AAG TTT ACA AAC AGT ATA CGG TTA CAC CCT GAT Val Phe Val Pro Ser Lys Phe Thr Asn Ser Ile Arg Leu His Pro Asp 465 470 475	1803
ACA AAT TAT GAT AAA ATC AAA GCC CAC CAT TGG ATA AAA ATT TGC CTT Thr Asn Tyr Asp Lys Ile Lys Ala His His Trp Ile Lys Ile Cys Leu 480 485 490	1851
CGT CTT TCC AAG AAG TAC GGG GAC AAT AGA AAA CAT TTC GAA ATA AGT Arg Leu Ser Lys Lys Tyr Gly Asp Asn Arg Lys His Phe Glu Ile Ser 495 500 505	1899
ATT GAT TCT CCA ATC CAT ATT TTA AAT CAA CTA TGC TCA CAC GCG AAT Ile Asp Ser Pro Ile His Ile Leu Asn Gln Leu Cys Ser His Ala Asn 510 515 520	1947
ACT TTG CTA CCG AGC TAC GAG AGT CAT TTC CAG TAT TGT GAT GAA GAT Thr Leu Leu Pro Ser Tyr Glu Ser His Phe Gln Tyr Cys Asp Glu Asp 525 530 535 540	1995
GGT AAT TTC GCA CCA GCA GCA GAT CAA CAA AAT TAC GCA AGT CAT CAT Gly Asn Phe Ala Pro Ala Ala Asp Gln Gln Asn Tyr Ala Ser His His 545 550 555	2043
GAT TCC AAT ATT TTC TTC CCA AAA GAA GTT CTT TCG TCT CCC GTT CTT Asp Ser Asn Ile Phe Phe Pro Lys Glu Val Leu Ser Ser Pro Val Leu 560 565 570	2091
TCA CCT AAC GTG CAG AAG ATG AAC ATT AGA ATA CCG TCT GAT CTT CCA Ser Pro Asn Val Gln Lys Met Asn Ile Arg Ile Pro Ser Asp Leu Pro 575 580 585	2139
GTA GTG CGT AAT AGA GCT GAA AGC GTA AAG AAA AGC AAG TCA GAT AAT Val Val Arg Asn Arg Ala Glu Ser Val Lys Lys Ser Lys Ser Asp Asn 590 595 600	2187

ACC	TCC	AAG	AAG	AAT	GAT	CAA	AGT	AGC	AAT	GTC	TTC	GCA	TCC	AAA	CAG	2235
Thr	Ser	Lys	Lys	Asn	Asp	Gln	Ser	Ser	Asn	Val	Phe	Ala	Ser	Lys	Gln	
605					610					615					620	
CTG	GTC	GCA	AAC	ATT	TAT	AAG	CCC	AAT	CAG	ATT	CCA	AGA	GAA	TTA	ACT	2283
Leu	Val	Ala	Asn	Ile	Tyr	Lys	Pro	Asn	Gln	Ile	Pro	Arg	Glu	Leu	Thr	
				625					630					635		
TCT	CCT	CAG	GCG	TTA	CCA	TTA	TCG	CCC	ATC	ACC	TCA	CCA	ATT	CTC	AAT	2331
Ser	Pro	Gln	Ala	Leu	Pro	Leu	Ser	Pro	Ile	Thr	Ser	Pro	Ile	Leu	Asn	
			640					645					650			
TAC	CAA	CCA	TTA	TCA	AAC	TCC	CCG	CCT	CCA	GAT	TTT	GAT	TTT	GAT	CTA	2379
Tyr	Gln	Pro	Leu	Ser	Asn	Ser	Pro	Pro	Pro	Asp	Phe	Asp	Phe	Asp	Leu	
		655					660					665				
GCT	AAG	CGC	GGC	GCA	GCC	GAT	TCT	CAT	GCT	ATT	CCT	GTG	GAT	CCT	CCA	2427
Ala	Lys	Arg	Gly	Ala	Ala	Asp	Ser	His	Ala	Ile	Pro	Val	Asp	Pro	Pro	
	670					675					680					
TCA	TAT	TTT	GAT	GTA	TTA	AAG	GCC	GAT	GGG	ATT	GAA	TTG	CCA	TAC	TAC	2475
Ser	Tyr	Phe	Asp	Val	Leu	Lys	Ala	Asp	Gly	Ile	Glu	Leu	Pro	Tyr	Tyr	
685					690					695					700	
GAT	ACA	AGT	TCA	TCT	AAA	ATT	CCT	GAA	CTA	AAA	CTA	AAC	AAA	TCT	AGA	2523
Asp	Thr	Ser	Ser	Ser	Lys	Ile	Pro	Glu	Leu	Lys	Leu	Asn	Lys	Ser	Arg	
				705					710					715		
GAG	ACA	TTG	GCC	AGC	ATT	GAG	GAG	GAC	TCA	TTC	AAT	GGT	TGG	TCT	CAA	2571
Glu	Thr	Leu	Ala	Ser	Ile	Glu	Glu	Asp	Ser	Phe	Asn	Gly	Trp	Ser	Gln	
			720					725					730			
ATT	GAT	GAC	TTA	TCC	GAC	GAA	GAT	GAC	AAT	GAT	GGC	GAT	ATA	GCA	TCT	2619
Ile	Asp	Asp	Leu	Ser	Asp	Glu	Asp	Asp	Asn	Asp	Gly	Asp	Ile	Ala	Ser	
		735					740					745				
GGT	TTC	AAC	TTC	AAG	CTG	TCA	ACC	AGT	GCT	CCG	AGT	GAG	AAC	GTT	AAT	2667
Gly	Phe	Asn	Phe	Lys	Leu	Ser	Thr	Ser	Ala	Pro	Ser	Glu	Asn	Val	Asn	
	750					755					760					
TCA	CAC	ACT	CCT	ATT	TTG	CAG	TCT	TTA	AAC	ATG	AGT	CTT	GAT	GGG	AGA	2715
Ser	His	Thr	Pro	Ile	Leu	Gln	Ser	Leu	Asn	Met	Ser	Leu	Asp	Gly	Arg	
765					770					775					780	
AAA	AAA	AAT	CGT	GCC	AGT	CTA	CAC	GCA	ACA	TCA	GTG	TTA	CCT	AGT	ACA	2763
Lys	Lys	Asn	Arg	Ala	Ser	Leu	His	Ala	Thr	Ser	Val	Leu	Pro	Ser	Thr	
				785					790					795		
ATA	AGA	CAG	AAC	AAT	CAG	CAT	TTC	AAT	GAC	ATA	AAC	CAG	ATG	CTA	GGC	2811
Ile	Arg	Gln	Asn	Asn	Gln	His	Phe	Asn	Asp	Ile	Asn	Gln	Met	Leu	Gly	
			800					805					810			
AGT	AGT	GAC	GAA	GAT	GCC	TTT	CCC	AAA	AGC	CAA	TCA	TTA	AAT	TTC	AAT	2859
Ser	Ser	Asp	Glu	Asp	Ala	Phe	Pro	Lys	Ser	Gln	Ser	Leu	Asn	Phe	Asn	
		815					820					825				
AAG	AAA	CTA	CCA	ATA	CTT	AAA	ATT	AAT	GAT	AAC	GTC	ATA	CAA	TCA	AAC	2907
Lys	Lys	Leu	Pro	Ile	Leu	Lys	Ile	Asn	Asp	Asn	Val	Ile	Gln	Ser	Asn	
		830				835					840					
AGC	AAT	AGT	AAT	AAC	AGA	GTT	GAT	AAT	CCA	GAA	GAT	ACA	GTG	GAT	TCT	2955
Ser	Asn	Ser	Asn	Asn	Arg	Val	Asp	Asn	Pro	Glu	Asp	Thr	Val	Asp	Ser	
845					850					855					860	

46

TCA GTC GAT ATT ACA GCA TTT TAT GAT CCA AGA ATG TCA TCA GAT TCC Ser Val Asp Ile Thr Ala Phe Tyr Asp Pro Arg Met Ser Ser Asp Ser 865 870 875	3003
AAA TTT GAT TGG GAG GTA AGC AAG AAC CAT GTT GAC CCA GCA GCC TAC Lys Phe Asp Trp Glu Val Ser Lys Asn His Val Asp Pro Ala Ala Tyr 880 885 890	3051
TCG GTT AAC GTT GCT AGT GAA AAC CGT GTA CTG GAC GAC TTT AAG AAA Ser Val Asn Val Ala Ser Glu Asn Arg Val Leu Asp Asp Phe Lys Lys 895 900 905	3099
GCA TTT CGC GAA AAG AGA AAA TAAGTACATT ATTTTCATTC TCCGACAGAA Ala Phe Arg Glu Lys Arg Lys 910 915	3150
TTGCTACCAT TTTACTTTGT GTCCTGTGAT TCAATAGTGT ACAATATATT GGACATTTTA	3210
TAGTATACAA ATATACACCA TCAATCTATA CATCCATATC ACTTGTCGTA AAGATATCCC	3270
TTTTTAATAG TACAGCGATT AAAAAAATAA CATGATTAAAC GTTCAGTTAC CAATGAGCTT	3330
ATTTATTAGG CTTGCTTTAG ATTTTTCCAA GTCAATTTTT GTTTTTTCTA ACGCTTGCAA	3390
CCTCATCTCA ACCTTCTTCC TTGCAAGCA GATCTTCGAA ACCATCTCGT TTATTCTCTC	3450
AATGCTGTTT CCACTTTCAT CATCGTCTGG GAAAGTACC GGTAAGGGCG	3500

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 915 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Leu Gln Phe Asn Thr Glu Asn Asp Thr Val Ala Pro Val Phe Pro 1 5 10 15
Met Glu Gln Asp Ile Asn Ala Ala Pro Asp Ala Val Pro Leu Val Gln 20 25 30
Thr Thr Thr Leu Gln Val Phe Val Lys Leu Ala Glu Pro Ile Val Phe 35 40 45
Leu Lys Gly Phe Glu Thr Asn Gly Leu Ser Glu Ile Ala Pro Ser Ile 50 55 60
Leu Arg Gly Ser Leu Ile Val Arg Val Leu Lys Pro Asn Lys Leu Lys 65 70 75 80
Ser Ile Ser Ile Thr Phe Lys Gly Ile Ser Arg Thr Glu Trp Pro Glu 85 90 95
Gly Ile Pro Pro Lys Arg Glu Glu Phe Ser Asp Val Glu Thr Val Val 100 105 110
Asn His Thr Trp Pro Phe Tyr Gln Ala Asp Asp Gly Met Asn Ser Phe 115 120 125
Thr Leu Glu His His Ser Ser Asn Asn Ser Ser Asn Arg Pro Ser Met 130 135 140



Ser Asp Glu Asp Tyr Leu Leu Glu Lys Ser Gly Ala Ser Val Tyr Ile  
 145 150 155 160  
 Pro Pro Thr Ala Glu Pro Pro Lys Asp Asn Ser Asn Leu Ser Leu Asp  
 165 170 175  
 Ala Tyr Glu Arg Asn Ser Leu Ser Ser Asp Asn Leu Ser Asn Lys Pro  
 180 185 190  
 Val Ser Ser Asp Val Ser His Asp Asp Ser Lys Leu Leu Ala Ile Gln  
 195 200 205  
 Lys Thr Pro Leu Pro Ser Ser Ser Arg Arg Gly Ser Val Pro Ala Asn  
 210 215 220  
 Phe His Gly Asn Ser Leu Ser Pro His Thr Phe Ile Ser Asp Leu Phe  
 225 230 235 240  
 Thr Lys Thr Phe Ser Asn Ser Gly Ala Thr Pro Ser Pro Glu Gln Glu  
 245 250 255  
 Asp Asn Tyr Leu Thr Pro Ser Lys Asp Ser Lys Glu Val Phe Ile Phe  
 260 265 270  
 Arg Pro Gly Asp Tyr Ile Tyr Thr Phe Glu Gln Pro Ile Ser Gln Ser  
 275 280 285  
 Tyr Pro Glu Ser Ile Lys Ala Asn Phe Gly Ser Val Glu Tyr Lys Leu  
 290 295 300  
 Ser Ile Asp Ile Glu Arg Phe Gly Ala Phe Lys Ser Thr Ile His Thr  
 305 310 315 320  
 Gln Leu Pro Ile Lys Val Val Arg Leu Pro Ser Asp Gly Ser Val Glu  
 325 330 335  
 Glu Thr Glu Ala Ile Ala Ile Ser Lys Asp Trp Lys Asp Leu Leu His  
 340 345 350  
 Tyr Asp Val Val Ile Phe Ser Lys Glu Ile Val Leu Asn Ala Phe Leu  
 355 360 365  
 Pro Ile Asp Phe His Phe Ala Pro Leu Asp Lys Val Thr Leu His Arg  
 370 375 380  
 Ile Arg Ile Tyr Leu Thr Glu Ser Met Glu Tyr Thr Cys Asn Ser Asn  
 385 390 395 400  
 Gly Asn His Glu Lys Ala Arg Arg Leu Glu Pro Thr Lys Lys Phe Leu  
 405 410 415  
 Leu Ala Glu His Asn Gly Pro Lys Leu Pro His Ile Pro Ala Gly Ser  
 420 425 430  
 Asn Pro Leu Lys Ala Lys Asn Arg Gly Asn Ile Leu Leu Asp Glu Lys  
 435 440 445  
 Ser Gly Asp Leu Val Asn Lys Asp Phe Gln Phe Glu Val Phe Val Pro  
 450 455 460  
 Ser Lys Phe Thr Asn Ser Ile Arg Leu His Pro Asp Thr Asn Tyr Asp  
 465 470 475 480  
 Lys Ile Lys Ala His Trp Ile Lys Ile Cys Leu Arg Leu Ser Lys  
 485 490 495

Lys Tyr Gly Asp Asn Arg Lys His Phe Glu Ile Ser Ile Asp Ser Pro  
 500 505 510  
 Ile His Ile Leu Asn Gln Leu Cys Ser His Ala Asn Thr Leu Leu Pro  
 515 520 525  
 Ser Tyr Glu Ser His Phe Gln Tyr Cys Asp Glu Asp Gly Asn Phe Ala  
 530 535 540  
 Pro Ala Ala Asp Gln Gln Asn Tyr Ala Ser His His Asp Ser Asn Ile  
 545 550 555 560  
 Phe Phe Pro Lys Glu Val Leu Ser Ser Pro Val Leu Ser Pro Asn Val  
 565 570 575  
 Gln Lys Met Asn Ile Arg Ile Pro Ser Asp Leu Pro Val Val Arg Asn  
 580 585 590  
 Arg Ala Glu Ser Val Lys Lys Ser Lys Ser Asp Asn Thr Ser Lys Lys  
 595 600 605  
 Asn Asp Gln Ser Ser Asn Val Phe Ala Ser Lys Gln Leu Val Ala Asn  
 610 615 620  
 Ile Tyr Lys Pro Asn Gln Ile Pro Arg Glu Leu Thr Ser Pro Gln Ala  
 625 630 635 640  
 Leu Pro Leu Ser Pro Ile Thr Ser Pro Ile Leu Asn Tyr Gln Pro Leu  
 645 650 655  
 Ser Asn Ser Pro Pro Pro Asp Phe Asp Phe Asp Leu Ala Lys Arg Gly  
 660 665 670  
 Ala Ala Asp Ser His Ala Ile Pro Val Asp Pro Pro Ser Tyr Phe Asp  
 675 680 685  
 Val Leu Lys Ala Asp Gly Ile Glu Leu Pro Tyr Tyr Asp Thr Ser Ser  
 690 695 700  
 Ser Lys Ile Pro Glu Leu Lys Leu Asn Lys Ser Arg Glu Thr Leu Ala  
 705 710 715 720  
 Ser Ile Glu Glu Asp Ser Phe Asn Gly Trp Ser Gln Ile Asp Asp Leu  
 725 730 735  
 Ser Asp Glu Asp Asp Asn Asp Gly Asp Ile Ala Ser Gly Phe Asn Phe  
 740 745 750  
 Lys Leu Ser Thr Ser Ala Pro Ser Glu Asn Val Asn Ser His Thr Pro  
 755 760 765  
 Ile Leu Gln Ser Leu Asn Met Ser Leu Asp Gly Arg Lys Lys Asn Arg  
 770 775 780  
 Ala Ser Leu His Ala Thr Ser Val Leu Pro Ser Thr Ile Arg Gln Asn  
 785 790 795 800  
 Asn Gln His Phe Asn Asp Ile Asn Gln Met Leu Gly Ser Ser Asp Glu  
 805 810 815  
 Asp Ala Phe Pro Lys Ser Gln Ser Leu Asn Phe Asn Lys Lys Leu Pro  
 820 825 830  
 Ile Leu Lys Ile Asn Asp Asn Val Ile Gln Ser Asn Ser Asn Ser Asn  
 835 840 845

Asn Arg Val Asp Asn Pro Glu Asp Thr Val Asp Ser Ser Val Asp Ile  
 850 855 860  
 Thr Ala Phe Tyr Asp Pro Arg Met Ser Ser Asp Ser Lys Phe Asp Trp  
 865 870 875 880  
 Glu Val Ser Lys Asn His Val Asp Pro Ala Ala Tyr Ser Val Asn Val  
 885 890 895  
 Ala Ser Glu Asn Arg Val Leu Asp Asp Phe Lys Lys Ala Phe Arg Glu  
 900 905 910  
 Lys Arg Lys  
 915

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2745 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: coding sequence of CNI

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ATGCTCCAAT TCAATACAGA AAATGATACT GTAGCTCCAG TGTTCCCAT GGAGCAAGAT	60
ATAAATGCAG CACCTGATGC CGTCCCCTG GTGCAGACAA CAACACTACA AGTCTTTGTA	120
AAGCTTGCCG AACCCATAGT GTTTTTTAAA GGATTTGAAA CTAACGGACT GTCTGAAATA	180
GCCCCCAGTA TCTTACGAGG ATCTCTTATC GTCAGGGTGT TGAAACCGAA TAAATTTAAA	240
AGTATATCGA TAACCTTCAA AGGAATATCC AGAACAGAGT GGCCGGAAGG TATACCACCG	300
AAGAGAGAAG AATTTTCAGA TGTTGAACT GTTGTCATC ACACATGGCC ATTTTATCAG	360
GCGGATGACG GCATGAATTC TTTCACCTTA GAACATCACA GCTCAAATAA TTCGTCCAAT	420
CGCCCATCTA TGAGCGATGA AGATTATCTA CTTGAAAAA GCGGTGCTTC AGTATATATC	480
CCACCAACCG CTGAACCCCC TAAAGATAAT AGCAATCTAA GTCTGGATGC CTATGAGCGC	540
AACTCATTGT CATCCGATAA TTTGAGTAAC AAGCCAGTAT CAAGTGATGT TTCCCATGAC	600
GACAGTAAAC TGTTGGCTAT TCAAAGACA CCATTACCAT CATCTAGTCG AAGAGGATCG	660
GTACCGGCAA ATTTTCACGG TAACTCTTG TCACCTCATA CCTTCATATC TGATTTGTTT	720
ACAAAAACAT TCAGTAATAG TGGCGCTACT CCAAGTCCTG AGCAAGAGGA TAACTATCTT	780
ACACCATCCA AAGATTCTAA AGAAGTTTTT ATTTTTCGAC CGGGCGATTA TATTTACACT	840
TTTGAACAGC CAATATCGCA ATCTTATCCA GAAAGTATAA AAGCCAATTT TGGTTCCGTG	900
GAGTATAAAC TGTCATAGA CATAGAGAGG TTTGGCGCAT TCAAATCAAC TATACATACT	960

CAATTACCCA	TCAAAGTCGT	AAGGCTTCCT	TCTGATGGAT	CCGTAGAAGA	GA CTGAAGCT	1020
ATTGCAATTT	CCAAGGACTG	GAAAGATCTT	CTTCATTATG	ACGTGGTAAT	TTTCTCGAAA	1080
GAGATCGTTT	TGAATGCATT	TTTACCCATC	GATTTCCATT	TCGCTCCTCT	AGATAAAGTT	1140
ACTCTGCATC	GTATTAGAAT	TTATCTAACA	GAGTCTATGG	AATACACTTG	TAATAGTAAT	1200
GGAAATCACG	AGAAGGCTCG	TAGATTAGAG	CCAATAAAA	AGTTTCTGTT	GGCTGAACAT	1260
AACGGTCCTA	AACTGCCTCA	TATACCAGCT	GGTTCGAATC	CTTTGAAGGC	TAAAAATAGA	1320
GGGAACATCC	TCTTGATGA	AAAATCCGGC	GATCTAGTTA	ACAAAGATT	TCAGTTCGAG	1380
GTGTTTGTCC	CAAGCAAGTT	TACAAACAGT	ATACGGTTAC	ACCCTGATAC	AAATTATGAT	1440
AAAATCAAAG	CCCACCATTG	GATAAAAATT	TGCCTTCGTC	TTTCCAAGAA	GTACGGGGAC	1500
AATAGAAAAC	ATTTCGAAAT	AAGTATTGAT	TCTCCAATCC	ATATTTTAAA	TCAACTATGC	1560
TCACACGCGA	ATACTTTGCT	ACCGAGCTAC	GAGAGTCATT	TCCAGTATTG	TGATGAAGAT	1620
GGTAATTTTCG	CACCAGCAGC	AGATCAACAA	AATTACGCAA	GTCATCATGA	TTCCAATATT	1680
TTCTTCCCAA	AAGAAGTTCT	TTCGTCTCCC	GTTCTTTCAC	CTAACGTGCA	GAAGATGAAC	1740
ATTAGAATAC	CGTCTGATCT	TCCAGTAGTG	CGTAATAGAG	CTGAAAGCGT	AAAGAAAAGC	1800
AAGTCAGATA	ATACCTCCAA	GAAGAATGAT	CAAAGTAGCA	ATGTCTTCGC	ATCCAAACAG	1860
CTGGTCGCAA	ACATTTATAA	GCCCAATCAG	ATTCCAAGAG	AATTAAC TTC	TCCTCAGGCG	1920
TTACCATTAT	CGCCCATCAC	CTCACCAATT	CTCAATTACC	AACCATTATC	AAACTCCCCG	1980
CCTCCAGATT	TTGATTTTGA	TCTAGCTAAG	CGCGGCGCAG	CCGATTCTCA	TGCTATTCTT	2040
GTGGATCCTC	CATCATATTT	TGATGTATTA	AAGGCCGATG	GGATTGAATT	GCCATACTAC	2100
GATACAAGTT	CATCTAAAT	TCCTGAAC TA	AACTAAACA	AATCTAGAGA	GACATTGGCC	2160
AGCATTGAGG	AGGACTCATT	CAATGGTTGG	TCTCAAATTG	ATGACTTATC	CGACGAAGAT	2220
GACAATGATG	GCGATATAGC	ATCTGGTTTC	AACTTCAAGC	TGTCAACCAG	TGCTCCGAGT	2280
GAGAACGTTA	ATTCACACAC	TCCTATTTTG	CAGTCTTTAA	ACATGAGTCT	TGATGGGAGA	2340
AAAAAAAATC	GTGCCAGTCT	ACACGCAACA	TCAGTGTTAC	CTAGTACAAT	AAGACAGAAC	2400
AATCAGCATT	TCAATGACAT	AAACCAGATG	CTAGGCAGTA	GTGACGAAGA	TGCCTTTCCC	2460
AAAAGCCAAT	CATTAAATTT	CAATAAGAAA	CTACCAATAC	TTAAAATTAA	TGATAACGTC	2520
ATACAATCAA	ACAGCAATAG	TAATAACAGA	GTTGATAATC	CAGAAGATAC	AGTGGATTCT	2580
TCAGTCGATA	TTACAGCATT	TTATGATCCA	AGAATGTCAT	CAGATTCCAA	ATTGATTGG	2640
GAGGTAAGCA	AGAACCATGT	TGACCCAGCA	GCCTACTCGG	TTAACGTTGC	TAGTGAAAAC	2700
CGTGTACTGG	ACGACTTTAA	GAAAGCATTT	CGCGAAAAGA	GAAAA		2745

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

51

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: CNI-PRC-A

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CCAAAAAAG AGATCTCGGA TCAAAGTAGC

30

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 30 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: CNI-PCR-B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GGGTTTTTCA GTGTCGACGA TTCATAGATC

30

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1964 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: genomic DNA fragment containing full  
CNA1 coding sequence

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 286..1944

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GTTTGTTGCA TTTTGATATT CATCTATATC TATTTCAAAA TTTTTCATGT CATCGCCTCT

60

TGAAACATGA	ATTTTCCAAT	TCTGAAAAAG	AACGTACTAC	TGGGAAACAA	AAGGGAAAAA	120
TGTATAAATC	CTTTAATGTT	TTTGAATCAA	GAGGCATTAT	TATAAAAGAA	CGAAGCAAAG	180
CCTTTAATAT	TTGCTTTATT	AAAGGTATTA	TTCAAAGAAA	AGTTTTTTTA	GATTCTTTTT	240
TTTTTGACGT	ATTAGCTCAG	CTGCCATAAA	ACACTCTCAA	CGCCA	ATG TCG AAA	294
					Met Ser Lys	
					1	
GAC TTG AAT TCT TCA CGC ATC AAA ATC ATT AAA CCT AAT GAC TCT TAC						342
Asp Leu Asn Ser Ser Arg Ile Lys Ile Ile Lys Pro Asn Asp Ser Tyr						
5 10 15						
ATA AAG GTT GAC CGG AAA AAA GAT TTA ACA AAA TAC GAA TTA GAA AAC						390
Ile Lys Val Asp Arg Lys Lys Asp Leu Thr Lys Tyr Glu Leu Glu Asn						
20 25 30 35						
GGT AAA GTA ATT TCT ACT AAG GAC CGA TCC TAC GCT TCT GTA CCT GCC						438
Gly Lys Val Ile Ser Thr Lys Asp Arg Ser Tyr Ala Ser Val Pro Ala						
40 45 50						
ATA ACA GGA AAG ATA CCA AGT GAT GAG GAA GTA TTC GAC TCC AAG ACG						486
Ile Thr Gly Lys Ile Pro Ser Asp Glu Glu Val Phe Asp Ser Lys Thr						
55 60 65						
GGA TTA CCT AAT CAT TCC TTT TTA AGA GAG CAT TTC TTT CAT GAG GGT						534
Gly Leu Pro Asn His Ser Phe Leu Arg Glu His Phe Phe His Glu Gly						
70 75 80						
CGA CTT TCT AAG GAA CAG GCC ATA AAA ATC TTA AAT ATG TCA ACT GTA						582
Arg Leu Ser Lys Glu Gln Ala Ile Lys Ile Leu Asn Met Ser Thr Val						
85 90 95						
GCA TTG AGT AAA GAA CCC AAT CTA CTA AAA CTC AAA GCG CCA ATT ACT						630
Ala Leu Ser Lys Glu Pro Asn Leu Leu Lys Leu Lys Ala Pro Ile Thr						
100 105 110 115						
ATA TGT GGT GAT ATT CAC GGG CAG TAT TAT GAT TTA TTG AAA CTG TTT						678
Ile Cys Gly Asp Ile His Gly Gln Tyr Tyr Asp Leu Leu Lys Leu Phe						
120 125 130						
GAA GTT GGC GGT GAC CCC GCC GAA ATC GAC TAT TTA TTC TTG GGG GAT						726
Glu Val Gly Gly Asp Pro Ala Glu Ile Asp Tyr Leu Phe Leu Gly Asp						
135 140 145						
TAT GTT GAT AGA GGT GCA TTC TCT TTT GAG TGT CTG ATT TAT TTG TAC						774
Tyr Val Asp Arg Gly Ala Phe Ser Phe Glu Cys Leu Ile Tyr Leu Tyr						
150 155 160						
TCC TTG AAG TTG AAT AAT TTA GGG AGA TTT TGG ATG CTA AGA GGT AAC						822
Ser Leu Lys Leu Asn Asn Leu Gly Arg Phe Trp Met Leu Arg Gly Asn						
165 170 175						
CAT GAG TGT AAG CAC TTG ACC TCT TAT TTT ACT TTT AAG AAT GAG ATG						870
His Glu Cys Lys His Leu Thr Ser Tyr Phe Thr Phe Lys Asn Glu Met						
180 185 190 195						
TTG CAC AAA TAC GAT ATG GAA GTT TAC GAT GCT TGC TGC AGA TCA TTC						918
Leu His Lys Tyr Asp Met Glu Val Tyr Asp Ala Cys Cys Arg Ser Phe						
200 205 210						
AAT GTC TTA CCA TTA GCA GCT TTA ATG AAC GGA CAA TAT TTT TGT GTG						966
Asn Val Leu Pro Leu Ala Ala Leu Met Asn Gly Gln Tyr Phe Cys Val						
215 220 225						

CAT	GGT	GGT	ATC	TCT	CCA	GAG	TTA	AAA	TCA	GTA	GAG	GAT	GTT	AAT	AAA	1014
His	Gly	Gly	Ile	Ser	Pro	Glu	Leu	Lys	Ser	Val	Glu	Asp	Val	Asn	Lys	
		230					235					240				
ATT	AAT	AGA	TTT	CGA	GAA	ATC	CCA	TCT	CGT	GGT	CTC	ATG	TGT	GAC	CTA	1062
Ile	Asn	Arg	Phe	Arg	Glu	Ile	Pro	Ser	Arg	Gly	Leu	Met	Cys	Asp	Leu	
	245					250					255					
CTA	TGG	GCC	GAT	CCT	GTC	GAA	AAT	TAT	GAT	GAT	GCA	AGA	GAT	GGT	AGC	1110
Leu	Trp	Ala	Asp	Pro	Val	Glu	Asn	Tyr	Asp	Asp	Ala	Arg	Asp	Gly	Ser	
260					265				270						275	
GAA	TTT	GAT	CAG	AGC	GAG	GAT	GAA	TTC	GTA	CCT	AAC	AGT	TTG	AGG	GGT	1158
Glu	Phe	Asp	Gln	Ser	Glu	Asp	Glu	Phe	Val	Pro	Asn	Ser	Leu	Arg	Gly	
				280					285					290		
TGC	TCT	TTC	GCC	TTC	ACT	TTT	AAA	GCA	TCA	TGC	AAG	TTT	TTG	AAG	GCA	1206
Cys	Ser	Phe	Ala	Phe	Thr	Phe	Lys	Ala	Ser	Cys	Lys	Phe	Leu	Lys	Ala	
			295					300					305			
AAT	GGT	TTG	TTA	TCT	ATT	ATT	AGA	GCA	CAC	GAA	GCA	CAG	GAT	GCT	GGG	1254
Asn	Gly	Leu	Leu	Ser	Ile	Ile	Arg	Ala	His	Glu	Ala	Gln	Asp	Ala	Gly	
		310					315					320				
TAC	AGA	ATG	TAT	AAA	AAC	AAT	AAA	GTA	ACA	GGC	TTC	CCG	AGC	TTA	ATA	1302
Tyr	Arg	Met	Tyr	Lys	Asn	Asn	Lys	Val	Thr	Gly	Phe	Pro	Ser	Leu	Ile	
	325					330					335					
ACC	ATG	TTC	AGT	GCG	CCA	AAC	TAC	CTG	GAC	ACA	TAT	CAT	AAT	AAA	GCT	1350
Thr	Met	Phe	Ser	Ala	Pro	Asn	Tyr	Leu	Asp	Thr	Tyr	His	Asn	Lys	Ala	
340					345					350					355	
GCT	GTG	TTA	AAA	TAT	GAA	GAA	AAC	GTC	ATG	AAC	ATC	AGG	CAG	TTT	CAC	1398
Ala	Val	Leu	Lys	Tyr	Glu	Glu	Asn	Val	Met	Asn	Ile	Arg	Gln	Phe	His	
				360					365					370		
ATG	TCT	CCG	CAC	CCT	TAC	TGG	TTG	CCT	GAT	TTT	ATG	GAT	GTT	TTC	ACC	1446
Met	Ser	Pro	His	Pro	Tyr	Trp	Leu	Pro	Asp	Phe	Met	Asp	Val	Phe	Thr	
			375					380					385			
TGG	TCA	CTA	CCT	TTT	GTT	GGC	GAA	AAA	GTT	ACT	AGC	ATG	TTA	GTG	TCT	1494
Trp	Ser	Leu	Pro	Phe	Val	Gly	Glu	Lys	Val	Thr	Ser	Met	Leu	Val	Ser	
		390					395					400				
ATA	TTA	AAC	ATA	TGT	AGT	GAG	CAG	GAA	CTT	GAC	CCA	GAA	TCG	GAA	CCC	1542
Ile	Leu	Asn	Ile	Cys	Ser	Glu	Gln	Glu	Leu	Asp	Pro	Glu	Ser	Glu	Pro	
	405					410					415					
AAA	GCT	GCG	GAG	GAG	ACT	GTA	AAG	GCA	AGA	GCA	AAC	GCA	ACT	AAG	GAG	1590
Lys	Ala	Ala	Glu	Glu	Thr	Val	Lys	Ala	Arg	Ala	Asn	Ala	Thr	Lys	Glu	
420					425				430						435	
ACC	GGC	ACC	CCA	TCT	GAT	GAA	AAG	GCG	TCA	TCA	GCG	ATA	TTA	GAA	GAT	1638
Thr	Gly	Thr	Pro	Ser	Asp	Glu	Lys	Ala	Ser	Ser	Ala	Ile	Leu	Glu	Asp	
				440					445					450		
GAA	ACC	CGA	AGA	AAG	GCT	TTG	AGA	AAT	AAG	ATA	TTA	GCT	ATT	GCT	AAA	1686
Glu	Thr	Arg	Arg	Lys	Ala	Leu	Arg	Asn	Lys	Ile	Leu	Ala	Ile	Ala	Lys	
				455				460					465			
GTT	TCA	AGA	ATG	TTT	TCG	GTG	CTA	AGA	GAA	GAG	AGC	GAA	AAA	GTG	GAA	1734
Val	Ser	Arg	Met	Phe	Ser	Val	Leu	Arg	Glu	Glu	Ser	Glu	Lys	Val	Glu	
		470					475					480				
TAT	TTG	AAA	ACT	ATG	AAT	GCC	GGT	GTC	TTA	CCT	CGT	GGT	GCT	CTA	GCT	1782
Tyr	Leu	Lys	Thr	Met	Asn	Ala	Gly	Val	Leu	Pro	Arg	Gly	Ala	Leu	Ala	
	485					490					495					

54

CGT GGG ACT GAA GGT TTG AAT GAA ACG CTA AGC ACT TTT GAA AAG GCT	1830
Arg Gly Thr Glu Gly Leu Asn Glu Thr Leu Ser Thr Phe Glu Lys Ala	
500 505 510 515	
AGA AAG GAA GAC CTT ATT AAT GAA AAA TTA CCA CCA TCT TTA TCG GAG	1878
Arg Lys Glu Asp Leu Ile Asn Glu Lys Leu Pro Pro Ser Leu Ser Glu	
520 525 530	
GTT GAA CAA GAG AAG ATT AAA TAC TAC GAA AAA ATA TTA AAG GGA GCG	1926
Val Glu Gln Glu Lys Ile Lys Tyr Tyr Glu Lys Ile Leu Lys Gly Ala	
535 540 545	
GAG AAA AAG CCA CAA CTG TGATAAATCT TCATTTTATT	1964
Glu Lys Lys Pro Gln Leu	
550	

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 553 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met	Ser	Lys	Asp	Leu	Asn	Ser	Ser	Arg	Ile	Lys	Ile	Ile	Lys	Pro	Asn	1	5	10	15
Asp	Ser	Tyr	Ile	Lys	Val	Asp	Arg	Lys	Lys	Asp	Leu	Thr	Lys	Tyr	Glu	20	25	30	
Leu	Glu	Asn	Gly	Lys	Val	Ile	Ser	Thr	Lys	Asp	Arg	Ser	Tyr	Ala	Ser	35	40	45	
Val	Pro	Ala	Ile	Thr	Gly	Lys	Ile	Pro	Ser	Asp	Glu	Glu	Val	Phe	Asp	50	55	60	
Ser	Lys	Thr	Gly	Leu	Pro	Asn	His	Ser	Phe	Leu	Arg	Glu	His	Phe	Phe	65	70	75	80
His	Glu	Gly	Arg	Leu	Ser	Lys	Glu	Gln	Ala	Ile	Lys	Ile	Leu	Asn	Met	85	90	95	
Ser	Thr	Val	Ala	Leu	Ser	Lys	Glu	Pro	Asn	Leu	Leu	Lys	Leu	Lys	Ala	100	105	110	
Pro	Ile	Thr	Ile	Cys	Gly	Asp	Ile	His	Gly	Gln	Tyr	Tyr	Asp	Leu	Leu	115	120	125	
Lys	Leu	Phe	Glu	Val	Gly	Gly	Asp	Pro	Ala	Glu	Ile	Asp	Tyr	Leu	Phe	130	135	140	
Leu	Gly	Asp	Tyr	Val	Asp	Arg	Gly	Ala	Phe	Ser	Phe	Glu	Cys	Leu	Ile	145	150	155	160
Tyr	Leu	Tyr	Ser	Leu	Lys	Leu	Asn	Asn	Leu	Gly	Arg	Phe	Trp	Met	Leu	165	170	175	
Arg	Gly	Asn	His	Glu	Cys	Lys	His	Leu	Thr	Ser	Tyr	Phe	Thr	Phe	Lys	180	185	190	
Asn	Glu	Met	Leu	His	Lys	Tyr	Asp	Met	Glu	Val	Tyr	Asp	Ala	Cys	Cys	195	200	205	



55

Arg Ser Phe Asn Val Leu Pro Leu Ala Ala Leu Met Asn Gly Gln Tyr  
 210 215 220  
 Phe Cys Val His Gly Gly Ile Ser Pro Glu Leu Lys Ser Val Glu Asp  
 225 230 235 240  
 Val Asn Lys Ile Asn Arg Phe Arg Glu Ile Pro Ser Arg Gly Leu Met  
 245 250 255  
 Cys Asp Leu Leu Trp Ala Asp Pro Val Glu Asn Tyr Asp Asp Ala Arg  
 260 265 270  
 Asp Gly Ser Glu Phe Asp Gln Ser Glu Asp Glu Phe Val Pro Asn Ser  
 275 280 285  
 Leu Arg Gly Cys Ser Phe Ala Phe Thr Phe Lys Ala Ser Cys Lys Phe  
 290 295 300  
 Leu Lys Ala Asn Gly Leu Leu Ser Ile Ile Arg Ala His Glu Ala Gln  
 305 310 315 320  
 Asp Ala Gly Tyr Arg Met Tyr Lys Asn Asn Lys Val Thr Gly Phe Pro  
 325 330 335  
 Ser Leu Ile Thr Met Phe Ser Ala Pro Asn Tyr Leu Asp Thr Tyr His  
 340 345 350  
 Asn Lys Ala Ala Val Leu Lys Tyr Glu Glu Asn Val Met Asn Ile Arg  
 355 360 365  
 Gln Phe His Met Ser Pro His Pro Tyr Trp Leu Pro Asp Phe Met Asp  
 370 375 380  
 Val Phe Thr Trp Ser Leu Pro Phe Val Gly Glu Lys Val Thr Ser Met  
 385 390 395 400  
 Leu Val Ser Ile Leu Asn Ile Cys Ser Glu Gln Glu Leu Asp Pro Glu  
 405 410 415  
 Ser Glu Pro Lys Ala Ala Glu Glu Thr Val Lys Ala Arg Ala Asn Ala  
 420 425 430  
 Thr Lys Glu Thr Gly Thr Pro Ser Asp Glu Lys Ala Ser Ser Ala Ile  
 435 440 445  
 Leu Glu Asp Glu Thr Arg Arg Lys Ala Leu Arg Asn Lys Ile Leu Ala  
 450 455 460  
 Ile Ala Lys Val Ser Arg Met Phe Ser Val Leu Arg Glu Glu Ser Glu  
 465 470 475 480  
 Lys Val Glu Tyr Leu Lys Thr Met Asn Ala Gly Val Leu Pro Arg Gly  
 485 490 495  
 Ala Leu Ala Arg Gly Thr Glu Gly Leu Asn Glu Thr Leu Ser Thr Phe  
 500 505 510  
 Glu Lys Ala Arg Lys Glu Asp Leu Ile Asn Glu Lys Leu Pro Pro Ser  
 515 520 525  
 Leu Ser Glu Val Glu Gln Glu Lys Ile Lys Tyr Tyr Glu Lys Ile Leu  
 530 535 540  
 Lys Gly Ala Glu Lys Lys Pro Gln Leu  
 545 550

## (2) INFORMATION FOR SEQ ID NO:11:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2353 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

- (C) INDIVIDUAL ISOLATE: genomic DNA fragment containing full CNA2 coding sequence

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 262..2073

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATAGTCTATA ATACGTTTGA TACAGCTAGA TATCGCTAGC GCCAACATTG TCCCCCTCTC	60
TTGATCAATG CTTTTTTTTCG GCCCGAGACA AATGAGAAAA TGTCTTAAAA ATACCTTTCA	120
TCAAGACTCC TATTTTTCCT TAGAAAAAAC ATATATCCAA CTGGAACAGT ATTAAGCCAA	180
TTGCTACGAT ACAAACAAAA GGAGATATTC CTTCCCTCCC ATAGAGTCAC ACAGGAGCCA	240
GTACTTCTTC TTGAACCCGC A ATG TCT TCA GAC GCT ATA AGA AAT ACT GAG	291
Met Ser Ser Asp Ala Ile Arg Asn Thr Glu	
1 5 10	
CAG ATA AAC GCC GCT ATT AAA ATT ATA GAA AAC AAA ACA GAG CGT CCG	339
Gln Ile Asn Ala Ala Ile Lys Ile Ile Glu Asn Lys Thr Glu Arg Pro	
15 20 25	
CAA TCG TCC ACA ACC CCT ATA GAT TCG AAG GCT AGT ACA GTT GCT GCT	387
Gln Ser Ser Thr Thr Pro Ile Asp Ser Lys Ala Ser Thr Val Ala Ala	
30 35 40	
GCT AAT TCC ACG GCC ACA GAA ACT TCC AGA GAC CTT ACA CAA TAT ACC	435
Ala Asn Ser Thr Ala Thr Glu Thr Ser Arg Asp Leu Thr Gln Tyr Thr	
45 50 55	
CTA GAT GAC GGA AGA GTC GTA TCG ACA AAC CGC AGA ATA ATG AAT AAA	483
Leu Asp Asp Gly Arg Val Val Ser Thr Asn Arg Arg Ile Met Asn Lys	
60 65 70	
GTG CCC GCC ATC ACG TCA CAT GTT CCT ACA GAT GAA GAG CTG TTC CAG	531
Val Pro Ala Ile Thr Ser His Val Pro Thr Asp Glu Glu Leu Phe Gln	
75 80 85 90	
CCC AAT GGG ATA CCT CGT CAC GAA TTC CTA AGA GAT CAT TTC AAG CGC	579
Pro Asn Gly Ile Pro Arg His Glu Phe Leu Arg Asp His Phe Lys Arg	
95 100 105	
GAG GGC AAA TTG TCG GCT GCG CAG GCG GCC AGG ATC GTT ACA CTT GCA	627
Glu Gly Lys Leu Ser Ala Ala Gln Ala Ala Arg Ile Val Thr Leu Ala	
110 115 120	
ACG GAA CTC TTC AGC AAA GAA CCC AAC CTT ATA TCT GTT CCC GCC CCA	675
Thr Glu Leu Phe Ser Lys Glu Pro Asn Leu Ile Ser Val Pro Ala Pro	
125 130 135	

ATC	ACA	GTT	TGC	GGT	GAT	ATC	CAT	GGC	CAG	TAC	TTT	GAC	CTT	TTG	AAG	723
Ile	Thr	Val	Cys	Gly	Asp	Ile	His	Gly	Gln	Tyr	Phe	Asp	Leu	Leu	Lys	
140						145					150					
CTA	TTC	GAA	GTT	GGC	GGA	GAT	CCG	GCC	ACT	ACA	TCG	TAT	TTG	TTC	TTG	771
Leu	Phe	Glu	Val	Gly	Gly	Asp	Pro	Ala	Thr	Thr	Ser	Tyr	Leu	Phe	Leu	
155					160					165					170	
GGA	GAC	TAT	GTC	GAC	AGA	GGG	TCC	TTT	TCG	TTT	GAG	TGT	CTT	ATT	TAT	819
Gly	Asp	Tyr	Val	Asp	Arg	Gly	Ser	Phe	Ser	Phe	Glu	Cys	Leu	Ile	Tyr	
				175					180					185		
TTA	TAT	TCT	TTG	AAG	CTG	AAT	TTT	AAC	GAC	CAT	TTC	TGG	CTA	CTG	AGG	867
Leu	Tyr	Ser	Leu	Lys	Leu	Asn	Phe	Asn	Asp	His	Phe	Trp	Leu	Leu	Arg	
			190					195					200			
GGT	AAC	CAC	GAA	TGT	AAG	CAT	CTA	ACG	TCA	TAT	TTC	ACT	TTC	AAA	AAT	915
Gly	Asn	His	Glu	Cys	Lys	His	Leu	Thr	Ser	Tyr	Phe	Thr	Phe	Lys	Asn	
		205					210					215				
GAA	ATG	CTG	CAC	AAG	TAC	AAT	CTA	GAT	ATT	TAC	GAG	AAA	TGC	TGC	GAA	963
Glu	Met	Leu	His	Lys	Tyr	Asn	Leu	Asp	Ile	Tyr	Glu	Lys	Cys	Cys	Glu	
	220					225					230					
TCG	TTT	AAC	AAC	TTG	CCC	CTG	GCT	GCG	TTA	ATG	AAC	GGA	CAG	TAT	CTT	1011
Ser	Phe	Asn	Asn	Leu	Pro	Leu	Ala	Ala	Leu	Met	Asn	Gly	Gln	Tyr	Leu	
235					240					245					250	
TGT	GTT	CAT	GGT	GGT	ATA	TCT	CCC	GAG	TTA	AAC	TCT	TTA	CAG	GAC	ATT	1059
Cys	Val	His	Gly	Gly	Ile	Ser	Pro	Glu	Leu	Asn	Ser	Leu	Gln	Asp	Ile	
				255					260					265		
AAC	AAC	CTA	AAT	AGA	TTC	AGG	GAG	ATT	CCC	TCT	CAT	GGC	CTG	ATG	TGT	1107
Asn	Asn	Leu	Asn	Arg	Phe	Arg	Glu	Ile	Pro	Ser	His	Gly	Leu	Met	Cys	
			270					275					280			
GAT	CTG	TTG	TGG	GCT	GAC	CCG	ATT	GAA	GAG	TAC	GAC	GAA	GTC	TTG	GAT	1155
Asp	Leu	Leu	Trp	Ala	Asp	Pro	Ile	Glu	Glu	Tyr	Asp	Glu	Val	Leu	Asp	
		285					290					295				
AAA	GAC	TTG	ACT	GAG	GAA	GAC	ATA	GTG	AAC	TCC	AAA	ACC	ATG	GTT	CCT	1203
Lys	Asp	Leu	Thr	Glu	Glu	Asp	Ile	Val	Asn	Ser	Lys	Thr	Met	Val	Pro	
	300					305					310					
CAT	CAT	GGC	AAG	ATG	GCA	CCT	TCA	AGG	GAT	ATG	TTT	GTC	CCA	AAC	TCA	1251
His	His	Gly	Lys	Met	Ala	Pro	Ser	Arg	Asp	Met	Phe	Val	Pro	Asn	Ser	
315					320					325					330	
GTA	AGG	GGC	TGT	TCA	TAT	GCC	TTC	ACG	TAT	CGT	GCT	GCG	TGC	CAT	TTT	1299
Val	Arg	Gly	Cys	Ser	Tyr	Ala	Phe	Thr	Tyr	Arg	Ala	Ala	Cys	His	Phe	
				335					340					345		
CTG	CAA	GAG	ACT	GGC	CTG	TTG	TCC	ATC	ATC	AGG	GCA	CAC	GAG	GCT	CAA	1347
Leu	Gln	Glu	Thr	Gly	Leu	Leu	Ser	Ile	Ile	Arg	Ala	His	Glu	Ala	Gln	
			350					355					360			
GAC	GCT	GGT	TAT	AGA	ATG	TAC	AAA	AAT	ACC	AAG	ACT	TTG	GGC	TTT	CCC	1395
Asp	Ala	Gly	Tyr	Arg	Met	Tyr	Lys	Asn	Thr	Lys	Thr	Leu	Gly	Phe	Pro	
		365					370					375				
TCT	CTT	TTG	ACC	CTT	TTC	AGT	GCG	CCT	AAC	TAC	TTG	GAC	ACC	TAC	AAT	1443
Ser	Leu	Leu	Thr	Leu	Phe	Ser	Ala	Pro	Asn	Tyr	Leu	Asp	Thr	Tyr	Asn	
		380				385					390					
AAT	AAG	GCT	GCC	ATA	TTG	AAA	TAC	GAA	AAT	AAT	GTT	ATG	AAT	ATC	AGA	1491
Asn	Lys	Ala	Ala	Ile	Leu	Lys	Tyr	Glu	Asn	Asn	Val	Met	Asn	Ile	Arg	
395					400					405					410	

58

CAA TTC AAC ATG ACT CCA CAC CCC TAT TGG TTA CCA GAT TTC ATG GAC Gln Phe Asn Met Thr Pro His Pro Tyr Trp Leu Pro Asp Phe Met Asp 415 420 425	1539
GTT TTC ACG TGG TCC TTG CCA TTT GTT GGT GAA AAA GTT ACA GAG ATG Val Phe Thr Trp Ser Leu Pro Phe Val Gly Glu Lys Val Thr Glu Met 430 435 440	1587
CTT GTC GCA ATT CTA AAC ATC TGT ACT GAA GAT GAG CTG GAA AAC GAC Leu Val Ala Ile Leu Asn Ile Cys Thr Glu Asp Glu Leu Glu Asn Asp 445 450 455	1635
ACC CCC GTC ATT GAA GAA TTA GTT GGT ACC GAT AAA AAA TTG CCA CAA Thr Pro Val Ile Glu Glu Leu Val Gly Thr Asp Lys Lys Leu Pro Gln 460 465 470	1683
GCT GGT AAG TCG GAA GCA ACT CCA CAA CCA GCC ACT TCG GCG TCG CCT Ala Gly Lys Ser Glu Ala Thr Pro Gln Pro Ala Thr Ser Ala Ser Pro 475 480 485 490	1731
AAA CAT GCT TCC ATT TTA GAT GAC GAA CAT CGA AGG AAA GCC TTA CGA Lys His Ala Ser Ile Leu Asp Asp Glu His Arg Arg Lys Ala Leu Arg 495 500 505	1779
AAT AAG ATT CTG GCC GTC GCC AAA GTT TCC AGA ATG TAT TCT GTT CTC Asn Lys Ile Leu Ala Val Ala Lys Val Ser Arg Met Tyr Ser Val Leu 510 515 520	1827
AGA GAA GAA ACC AAT AAA GTT CAG TTT TTA AAA GAT CAC AAT TCA GGC Arg Glu Glu Thr Asn Lys Val Gln Phe Leu Lys Asp His Asn Ser Gly 525 530 535	1875
GTG TTG CCA CGT GGC GCT TTA TCT AAT GGT GTA AAG GGT TTA GAT GAA Val Leu Pro Arg Gly Ala Leu Ser Asn Gly Val Lys Gly Leu Asp Glu 540 545 550	1923
GCC CTG TCT ACC TTT GAA AGG GCA AGA AAG CAC GAT TTA ATT AAT GAA Ala Leu Ser Thr Phe Glu Arg Ala Arg Lys His Asp Leu Ile Asn Glu 555 560 565 570	1971
AAA TTA CCG CCT TCA CTA GAC GAA CTG AAA AAC GAA AAT AAG AAG TAC Lys Leu Pro Pro Ser Leu Asp Glu Leu Lys Asn Glu Asn Lys Lys Tyr 575 580 585	2019
TAC GAA AAA GTT TGG CAG AAA GTA CAT GAA CAT GAT GCA AAG AAT GAT Tyr Glu Lys Val Trp Gln Lys Val His Glu His Asp Ala Lys Asn Asp 590 595 600	2067
AGC AAA TAGAGAAAGC TCCTATTTCC ACTGTACATA CTTCAATAAG TAAGTAAGTT Ser Lys	2123
GCATTAATTA TCTATTTAGA AGCTAGATGC TCCTCAAATG CACAGAATCA TATAGCGTTT	2183
TATTAGGTCT GTTCTTTATT TTAGTTTTGT TGATCTCTAT GAAGGTATAT TTATATGCAA	2243
AAATAAACTT TTAAATATCT ATGGATGCTT ACTCAATTGT ATAGACGTTT TTCATAGGAG	2303
TGCAAATTAT GGACACCACC TTCTAATTGA GCAGAAGCGG TTCTGAATTC	2353

## (2) INFORMATION FOR SEQ ID NO:12:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 604 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

```

Met Ser Ser Asp Ala Ile Arg Asn Thr Glu Gln Ile Asn Ala Ala Ile
 1           5           10           15
Lys Ile Ile Glu Asn Lys Thr Glu Arg Pro Gln Ser Ser Thr Thr Pro
      20           25           30
Ile Asp Ser Lys Ala Ser Thr Val Ala Ala Ala Asn Ser Thr Ala Thr
      35           40           45
Glu Thr Ser Arg Asp Leu Thr Gln Tyr Thr Leu Asp Asp Gly Arg Val
      50           55           60
Val Ser Thr Asn Arg Arg Ile Met Asn Lys Val Pro Ala Ile Thr Ser
      65           70           75           80
His Val Pro Thr Asp Glu Glu Leu Phe Gln Pro Asn Gly Ile Pro Arg
      85           90           95
His Glu Phe Leu Arg Asp His Phe Lys Arg Glu Gly Lys Leu Ser Ala
      100          105          110
Ala Gln Ala Ala Arg Ile Val Thr Leu Ala Thr Glu Leu Phe Ser Lys
      115          120          125
Glu Pro Asn Leu Ile Ser Val Pro Ala Pro Ile Thr Val Cys Gly Asp
      130          135          140
Ile His Gly Gln Tyr Phe Asp Leu Leu Lys Leu Phe Glu Val Gly Gly
      145          150          155          160
Asp Pro Ala Thr Thr Ser Tyr Leu Phe Leu Gly Asp Tyr Val Asp Arg
      165          170          175
Gly Ser Phe Ser Phe Glu Cys Leu Ile Tyr Leu Tyr Ser Leu Lys Leu
      180          185          190
Asn Phe Asn Asp His Phe Trp Leu Leu Arg Gly Asn His Glu Cys Lys
      195          200          205
His Leu Thr Ser Tyr Phe Thr Phe Lys Asn Glu Met Leu His Lys Tyr
      210          215          220
Asn Leu Asp Ile Tyr Glu Lys Cys Cys Glu Ser Phe Asn Asn Leu Pro
      225          230          235          240
Leu Ala Ala Leu Met Asn Gly Gln Tyr Leu Cys Val His Gly Gly Ile
      245          250          255
Ser Pro Glu Leu Asn Ser Leu Gln Asp Ile Asn Asn Leu Asn Arg Phe
      260          265          270
Arg Glu Ile Pro Ser His Gly Leu Met Cys Asp Leu Leu Trp Ala Asp
      275          280          285
Pro Ile Glu Glu Tyr Asp Glu Val Leu Asp Lys Asp Leu Thr Glu Glu
      290          295          300
Asp Ile Val Asn Ser Lys Thr Met Val Pro His His Gly Lys Met Ala
      305          310          315          320
Pro Ser Arg Asp Met Phe Val Pro Asn Ser Val Arg Gly Cys Ser Tyr
      325          330          335

```

60

Ala Phe Thr Tyr Arg Ala Ala Cys His Phe Leu Gln Glu Thr Gly Leu  
                   340                  345                  350  
 Leu Ser Ile Ile Arg Ala His Glu Ala Gln Asp Ala Gly Tyr Arg Met  
                   355                  360                  365  
 Tyr Lys Asn Thr Lys Thr Leu Gly Phe Pro Ser Leu Leu Thr Leu Phe  
                   370                  375                  380  
 Ser Ala Pro Asn Tyr Leu Asp Thr Tyr Asn Asn Lys Ala Ala Ile Leu  
                   385                  390                  395                  400  
 Lys Tyr Glu Asn Asn Val Met Asn Ile Arg Gln Phe Asn Met Thr Pro  
                   405                  410                  415  
 His Pro Tyr Trp Leu Pro Asp Phe Met Asp Val Phe Thr Trp Ser Leu  
                   420                  425                  430  
 Pro Phe Val Gly Glu Lys Val Thr Glu Met Leu Val Ala Ile Leu Asn  
                   435                  440                  445  
 Ile Cys Thr Glu Asp Glu Leu Glu Asn Asp Thr Pro Val Ile Glu Glu  
                   450                  455                  460  
 Leu Val Gly Thr Asp Lys Lys Leu Pro Gln Ala Gly Lys Ser Glu Ala  
                   465                  470                  475                  480  
 Thr Pro Gln Pro Ala Thr Ser Ala Ser Pro Lys His Ala Ser Ile Leu  
                   485                  490                  495  
 Asp Asp Glu His Arg Arg Lys Ala Leu Arg Asn Lys Ile Leu Ala Val  
                   500                  505                  510  
 Ala Lys Val Ser Arg Met Tyr Ser Val Leu Arg Glu Glu Thr Asn Lys  
                   515                  520                  525  
 Val Gln Phe Leu Lys Asp His Asn Ser Gly Val Leu Pro Arg Gly Ala  
                   530                  535                  540  
 Leu Ser Asn Gly Val Lys Gly Leu Asp Glu Ala Leu Ser Thr Phe Glu  
                   545                  550                  555                  560  
 Arg Ala Arg Lys His Asp Leu Ile Asn Glu Lys Leu Pro Pro Ser Leu  
                   565                  570                  575  
 Asp Glu Leu Lys Asn Glu Asn Lys Lys Tyr Tyr Glu Lys Val Trp Gln  
                   580                  585                  590  
 Lys Val His Glu His Asp Ala Lys Asn Asp Ser Lys  
                   595                  600

## (2) INFORMATION FOR SEQ ID NO:13:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 812 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA (genomic)

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (C) INDIVIDUAL ISOLATE: genomic DNA fragment containing full

## CNB1 coding sequence

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 54..104

## (ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 181..652

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

ACTTGGTAAC TCAATGGTGA TCAGAATCCA TAGAAGCATT TTTATTCTT AAA ATG	56
Met	
1	
GGT GCT GCT CCT TCC AAA ATT GTG GAT GGT CTT TTA GAA GAT ACA AAT	104
Gly Ala Ala Pro Ser Lys Ile Val Asp Gly Leu Leu Glu Asp Thr Asn	
5 10 15	
TGTATGTACA CTTCTGGAGTG AGGAAAAGAA AGAAAGGGGA AATTAACCGA TTTACTAAC	164
ACTGACACTT TGAACA GTT GAT AGA GAT GAA ATT GAA AGG TTA AGG AAG	213
Val Asp Arg Asp Glu Ile Glu Arg Leu Arg Lys	
1 5 10	
AGA TTC ATG AAA TTA GAT AGA GAT AGC TCA GGG TCT ATT GAT AAA AAT	261
Arg Phe Met Lys Leu Asp Arg Asp Ser Ser Gly Ser Ile Asp Lys Asn	
15 20 25	
GAA TTT ATG AGC ATT CCT GGC GTT TCG TCA AAC CCT CTT GCT GGA CGT	309
Glu Phe Met Ser Ile Pro Gly Val Ser Ser Asn Pro Leu Ala Gly Arg	
30 35 40	
ATA ATG GAG GTT TTC GAT GCT GAT AAT AGT GGG GAC GTG GAT TTT CAA	357
Ile Met Glu Val Phe Asp Ala Asp Asn Ser Gly Asp Val Asp Phe Gln	
45 50 55	
GAG TTC ATC ACA GGA TTA TCC ATT TTC AGT GGG CGT GGG TCC AAG GAC	405
Glu Phe Ile Thr Gly Leu Ser Ile Phe Ser Gly Arg Gly Ser Lys Asp	
60 65 70 75	
GAA AAG TTA AGA TTC GCC TTC AAA ATC TAC GAC ATT GAC AAG GAC GGT	453
Glu Lys Leu Arg Phe Ala Phe Lys Ile Tyr Asp Ile Asp Lys Asp Gly	
80 85 90	
TTC ATA TCC AAT GGT GAG TTG TTC ATC GTG TTG AAG ATT ATG GTA GGT	501
Phe Ile Ser Asn Gly Glu Leu Phe Ile Val Leu Lys Ile Met Val Gly	
95 100 105	
TCT AAT CTG GAC GAT GAA CAG CTG CAA CAG ATA GTA GAT AGG ACG ATA	549
Ser Asn Leu Asp Asp Glu Gln Leu Gln Gln Ile Val Asp Arg Thr Ile	
110 115 120	
GTG GAA AAC GAT AGC GAC GGC GAC GGA CGT TTA AGT TTC GAG GAG TTT	597
Val Glu Asn Asp Ser Asp Gly Asp Gly Arg Leu Ser Phe Glu Glu Phe	
125 130 135	
AAG AAT GCT ATC GAA ACC ACA GAA GTG GCC AAG AGT CTG ACA TTG CAA	645
Lys Asn Ala Ile Glu Thr Thr Glu Val Ala Lys Ser Leu Thr Leu Gln	
140 145 150 155	
TAC GATGTGTAAG ACTAGGGGAC ACTTCATTCA TTTATGGTAT GCCAATATTT	698
Tyr Asp	

TTAAGAAAAG AAGAATAATA CGCGATATTG TTTTAAAGG AAGGAACGCA CACTCGCCCA 758  
 GTTAGAGTGC TGATGATATA TACATATATA TATGTATATG TAACAAACAA TAAG 812

## (2) INFORMATION FOR SEQ ID NO:14:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met Gly Ala Ala Pro Ser Lys Ile Val Asp Gly Leu Leu Glu Asp Thr  
 1 5 10 15  
 Asn

## (2) INFORMATION FOR SEQ ID NO:15:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 157 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Val Asp Arg Asp Glu Ile Glu Arg Leu Arg Lys Arg Phe Met Lys Leu  
 1 5 10 15  
 Asp Arg Asp Ser Ser Gly Ser Ile Asp Lys Asn Glu Phe Met Ser Ile  
 20 25 30  
 Pro Gly Val Ser Ser Asn Pro Leu Ala Gly Arg Ile Met Glu Val Phe  
 35 40 45  
 Asp Ala Asp Asn Ser Gly Asp Val Asp Phe Gln Glu Phe Ile Thr Gly  
 50 55 60  
 Leu Ser Ile Phe Ser Gly Arg Gly Ser Lys Asp Glu Lys Leu Arg Phe  
 65 70 75 80  
 Ala Phe Lys Ile Tyr Asp Ile Asp Lys Asp Gly Phe Ile Ser Asn Gly  
 85 90 95  
 Glu Leu Phe Ile Val Leu Lys Ile Met Val Gly Ser Asn Leu Asp Asp  
 100 105 110  
 Glu Gln Leu Gln Gln Ile Val Asp Arg Thr Ile Val Glu Asn Asp Ser  
 115 120 125  
 Asp Gly Asp Gly Arg Leu Ser Phe Glu Glu Phe Lys Asn Ala Ile Glu  
 130 135 140  
 Thr Thr Glu Val Ala Lys Ser Leu Thr Leu Gln Tyr Asp  
 145 150 155



## (2) INFORMATION FOR SEQ ID NO:16:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 524 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: coding sequence of CNB1

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..524

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

ATG GGT GCT GCT CCT TCC AAA ATT GTG GAT GGT CTT TTA GAA GAT ACA	48
Met Gly Ala Ala Pro Ser Lys Ile Val Asp Gly Leu Leu Glu Asp Thr	
1 5 10 15	
AAT TTT GAT AGA GAT GAA ATT GAA AGG TTA AGG AAG AGA TTC ATG AAA	96
Asn Phe Asp Arg Asp Glu Ile Glu Arg Leu Arg Lys Arg Phe Met Lys	
20 25 30	
TTA GAT AGA GAT AGC TCA GGG TCT ATT GAT AAA AAT GAA TTT ATG AGC	144
Leu Asp Arg Asp Ser Ser Gly Ser Ile Asp Lys Asn Glu Phe Met Ser	
35 40 45	
ATT CCT GGC GTT TCG TCA AAC CCT CTT GCT GGA CGT ATA ATG GAG GTT	192
Ile Pro Gly Val Ser Ser Asn Pro Leu Ala Gly Arg Ile Met Glu Val	
50 55 60	
TTC GAT GCT GAT AAT AGT GGG GAC GTG GAT TTT CAA GAG TTC ATC ACA	240
Phe Asp Ala Asp Asn Ser Gly Asp Val Asp Phe Gln Glu Phe Ile Thr	
65 70 75 80	
GGA TTA TCC ATT TTC AGT GGG CGT GGG TCC AAG GAC GAA AAG TTA AGA	288
Gly Leu Ser Ile Phe Ser Gly Arg Gly Ser Lys Asp Glu Lys Leu Arg	
85 90 95	
TTC GCC TTC AAA ATC TAC GAC ATT GAC AAG GAC GGT TTC ATA TCC AAT	336
Phe Ala Phe Lys Ile Tyr Asp Ile Asp Lys Asp Gly Phe Ile Ser Asn	
100 105 110	
GGT GAG TTG TTC ATC GTG TTG AAG ATT ATG GTA GGT TCT AAT CTG GAC	384
Gly Glu Leu Phe Ile Val Leu Lys Ile Met Val Gly Ser Asn Leu Asp	
115 120 125	
GAT GAA CAG CTG CAA CAG ATA GTA GAT AGG ACG ATA GTG GAA AAC GAT	432
Asp Glu Gln Leu Gln Gln Ile Val Asp Arg Thr Ile Val Glu Asn Asp	
130 135 140	
AGC GAC GGC GAC GGA CGT TTA AGT TTC GAG GAG TTT AAG AAT GCT ATC	480
Ser Asp Gly Asp Gly Arg Leu Ser Phe Glu Glu Phe Lys Asn Ala Ile	
145 150 155 160	
GAA ACC ACA GAA GTG GCC AAG AGT CTG ACA TTG CAA TAC GAT GT	524
Glu Thr Thr Glu Val Ala Lys Ser Leu Thr Leu Gln Tyr Asp	
165 170	

## (2) INFORMATION FOR SEQ ID NO:17:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 174 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

```

Met Gly Ala Ala Pro Ser Lys Ile Val Asp Gly Leu Leu Glu Asp Thr
 1           5           10           15
Asn Phe Asp Arg Asp Glu Ile Glu Arg Leu Arg Lys Arg Phe Met Lys
          20           25           30
Leu Asp Arg Asp Ser Ser Gly Ser Ile Asp Lys Asn Glu Phe Met Ser
          35           40           45
Ile Pro Gly Val Ser Ser Asn Pro Leu Ala Gly Arg Ile Met Glu Val
          50           55           60
Phe Asp Ala Asp Asn Ser Gly Asp Val Asp Phe Gln Glu Phe Ile Thr
          65           70           75           80
Gly Leu Ser Ile Phe Ser Gly Arg Gly Ser Lys Asp Glu Lys Leu Arg
          85           90           95
Phe Ala Phe Lys Ile Tyr Asp Ile Asp Lys Asp Gly Phe Ile Ser Asn
          100          105          110
Gly Glu Leu Phe Ile Val Leu Lys Ile Met Val Gly Ser Asn Leu Asp
          115          120          125
Asp Glu Gln Leu Gln Gln Ile Val Asp Arg Thr Ile Val Glu Asn Asp
          130          135          140
Ser Asp Gly Asp Gly Arg Leu Ser Phe Glu Glu Phe Lys Asn Ala Ile
          145          150          155          160
Glu Thr Thr Glu Val Ala Lys Ser Leu Thr Leu Gln Tyr Asp
          165          170

```

## (2) INFORMATION FOR SEQ ID NO:18:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1812 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (C) INDIVIDUAL ISOLATE: DNA fragment containing CNAIdeltaC coding sequence

## (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 286..1812

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

GTTTGTGCA TTTTGATATT CATCTATATC TATTTCAAAA TTTTTCATGT CATCGCCTCT	60
TGAAACATGA ATTTTCCAAT TCTGAAAAAG AACGTACTAC TGGGAAACAA AAGGGAAAAA	120
TGTATAAATC CTTTAATGTT TTTGAATCAA GAGGCATTAT TATAAAAGAA CGAAGCAAAG	180
CCTTTAATAT TTGCTTTATT AAAGGTATTA TTCAAAGAAA AGTTTTTTTA GATTCTTTTT	240
TTTTTGACGT ATTAGCTCAG CTGCCATAAA ACACTCTCAA CGCCA ATG TCG AAA	294
Met Ser Lys	
1	
GAC TTG AAT TCT TCA CGC ATC AAA ATC ATT AAA CCT AAT GAC TCT TAC	342
Asp Leu Asn Ser Ser Arg Ile Lys Ile Ile Lys Pro Asn Asp Ser Tyr	
5 10 15	
ATA AAG GTT GAC CGG AAA AAA GAT TTA ACA AAA TAC GAA TTA GAA AAC	390
Ile Lys Val Asp Arg Lys Lys Asp Leu Thr Lys Tyr Glu Leu Glu Asn	
20 25 30 35	
GGT AAA GTA ATT TCT ACT AAG GAC CGA TCC TAC GCT TCT GTA CCT GCC	438
Gly Lys Val Ile Ser Thr Lys Asp Arg Ser Tyr Ala Ser Val Pro Ala	
40 45 50	
ATA ACA GGA AAG ATA CCA AGT GAT GAG GAA GTA TTC GAC TCC AAG ACG	486
Ile Thr Gly Lys Ile Pro Ser Asp Glu Glu Val Phe Asp Ser Lys Thr	
55 60 65	
GGA TTA CCT AAT CAT TCC TTT TTA AGA GAG CAT TTC TTT CAT GAG GGT	534
Gly Leu Pro Asn His Ser Phe Leu Arg Glu His Phe Phe His Glu Gly	
70 75 80	
CGA CTT TCT AAG GAA CAG GCC ATA AAA ATC TTA AAT ATG TCA ACT GTA	582
Arg Leu Ser Lys Glu Gln Ala Ile Lys Ile Leu Asn Met Ser Thr Val	
85 90 95	
GCA TTG AGT AAA GAA CCC AAT CTA CTA AAA CTC AAA GCG CCA ATT ACT	630
Ala Leu Ser Lys Glu Pro Asn Leu Leu Lys Leu Lys Ala Pro Ile Thr	
100 105 110 115	
ATA TGT GGT GAT ATT CAC GGG CAG TAT TAT GAT TTA TTG AAA CTG TTT	678
Ile Cys Gly Asp Ile His Gly Gln Tyr Tyr Asp Leu Leu Lys Leu Phe	
120 125 130	
GAA GTT GGC GGT GAC CCC GCC GAA ATC GAC TAT TTA TTC TTG GGG GAT	726
Glu Val Gly Gly Asp Pro Ala Glu Ile Asp Tyr Leu Phe Leu Gly Asp	
135 140 145	
TAT GTT GAT AGA GGT GCA TTC TCT TTT GAG TGT CTG ATT TAT TTG TAC	774
Tyr Val Asp Arg Gly Ala Phe Ser Phe Glu Cys Leu Ile Tyr Leu Tyr	
150 155 160	
TCC TTG AAG TTG AAT AAT TTA GGG AGA TTT TGG ATG CTA AGA GGT AAC	822
Ser Leu Lys Leu Asn Asn Leu Gly Arg Phe Trp Met Leu Arg Gly Asn	
165 170 175	
CAT GAG TGT AAG CAC TTG ACC TCT TAT TTT ACT TTT AAG AAT GAG ATG	870
His Glu Cys Lys His Leu Thr Ser Tyr Phe Thr Phe Lys Asn Glu Met	
180 185 190 195	
TTG CAC AAA TAC GAT ATG GAA GTT TAC GAT GCT TGC TGC AGA TCA TTC	918
Leu His Lys Tyr Asp Met Glu Val Tyr Asp Ala Cys Cys Arg Ser Phe	
200 205 210	

AAT GTC TTA CCA TTA GCA GCT TTA ATG AAC GGA CAA TAT TTT TGT GTG Asn Val Leu Pro Leu Ala Ala Leu Met Asn Gly Gln Tyr Phe Cys Val 215 220 225	966
CAT GGT GGT ATC TCT CCA GAG TTA AAA TCA GTA GAG GAT GTT AAT AAA His Gly Gly Ile Ser Pro Glu Leu Lys Ser Val Glu Asp Val Asn Lys 230 235 240	1014
ATT AAT AGA TTT CGA GAA ATC CCA TCT CGT GGT CTC ATG TGT GAC CTA Ile Asn Arg Phe Arg Glu Ile Pro Ser Arg Gly Leu Met Cys Asp Leu 245 250 255	1062
CTA TGG GCC GAT CCT GTC GAA AAT TAT GAT GAT GCA AGA GAT GGT AGC Leu Trp Ala Asp Pro Val Glu Asn Tyr Asp Asp Ala Arg Asp Gly Ser 260 265 270 275	1110
GAA TTT GAT CAG AGC GAG GAT GAA TTC GTA CCT AAC AGT TTG AGG GGT Glu Phe Asp Gln Ser Glu Asp Glu Phe Val Pro Asn Ser Leu Arg Gly 280 285 290	1158
TGC TCT TTC GCC TTC ACT TTT AAA GCA TCA TGC AAG TTT TTG AAG GCA Cys Ser Phe Ala Phe Thr Phe Lys Ala Ser Cys Lys Phe Leu Lys Ala 295 300 305	1206
AAT GGT TTG TTA TCT ATT ATT AGA GCA CAC GAA GCA CAG GAT GCT GGG Asn Gly Leu Leu Ser Ile Ile Arg Ala His Glu Ala Gln Asp Ala Gly 310 315 320	1254
TAC AGA ATG TAT AAA AAC AAT AAA GTA ACA GGC TTC CCG AGC TTA ATA Tyr Arg Met Tyr Lys Asn Asn Lys Val Thr Gly Phe Pro Ser Leu Ile 325 330 335	1302
ACC ATG TTC AGT GCG CCA AAC TAC CTG GAC ACA TAT CAT AAT AAA GCT Thr Met Phe Ser Ala Pro Asn Tyr Leu Asp Thr Tyr His Asn Lys Ala 340 345 350 355	1350
GCT GTG TTA AAA TAT GAA GAA AAC GTC ATG AAC ATC AGG CAG TTT CAC Ala Val Leu Lys Tyr Glu Glu Asn Val Met Asn Ile Arg Gln Phe His 360 365 370	1398
ATG TCT CCG CAC CCT TAC TGG TTG CCT GAT TTT ATG GAT GTT TTC ACC Met Ser Pro His Pro Tyr Trp Leu Pro Asp Phe Met Asp Val Phe Thr 375 380 385	1446
TGG TCA CTA CCT TTT GTT GGC GAA AAA GTT ACT AGC ATG TTA GTG TCT Trp Ser Leu Pro Phe Val Gly Glu Lys Val Thr Ser Met Leu Val Ser 390 395 400	1494
ATA TTA AAC ATA TGT AGT GAG CAG GAA CTT GAC CCA GAA TCG GAA CCC Ile Leu Asn Ile Cys Ser Glu Gln Glu Leu Asp Pro Glu Ser Glu Pro 405 410 415	1542
AAA GCT GCG GAG GAG ACT GTA AAG GCA AGA GCA AAC GCA ACT AAG GAG Lys Ala Ala Glu Glu Thr Val Lys Ala Arg Ala Asn Ala Thr Lys Glu 420 425 430 435	1590
ACC GGC ACC CCA TCT GAT GAA AAG GCG TCA TCA GCG ATA TTA GAA GAT Thr Gly Thr Pro Ser Asp Glu Lys Ala Ser Ser Ala Ile Leu Glu Asp 440 445 450	1638
GAA ACC CGA AGA AAG GCT TTG AGA AAT AAG ATA TTA GCT ATT GCT AAA Glu Thr Arg Arg Lys Ala Leu Arg Asn Lys Ile Leu Ala Ile Ala Lys 455 460 465	1686
GTT TCA AGA ATG TTT TCG GTG CTA AGA GAA GAG AGC GAA AAA GTG GAA Val Ser Arg Met Phe Ser Val Leu Arg Glu Glu Ser Glu Lys Val Glu 470 475 480	1734

67

TAT TTG AAA ACT ATG AAT GCC GGT GTC TTA CCT CGT GGT GCT CTA GCT 1782  
 Tyr Leu Lys Thr Met Asn Ala Gly Val Leu Pro Arg Gly Ala Leu Ala  
 485 490 495

CGT GGG ACT GAA GGT TTG AAT GAA ACG CTA 1812  
 Arg Gly Thr Glu Gly Leu Asn Glu Thr Leu  
 500 505

## (2) INFORMATION FOR SEQ ID NO:19:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 509 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Met Ser Lys Asp Leu Asn Ser Ser Arg Ile Lys Ile Ile Lys Pro Asn  
 1 5 10 15  
 Asp Ser Tyr Ile Lys Val Asp Arg Lys Lys Asp Leu Thr Lys Tyr Glu  
 20 25 30  
 Leu Glu Asn Gly Lys Val Ile Ser Thr Lys Asp Arg Ser Tyr Ala Ser  
 35 40 45  
 Val Pro Ala Ile Thr Gly Lys Ile Pro Ser Asp Glu Glu Val Phe Asp  
 50 55 60  
 Ser Lys Thr Gly Leu Pro Asn His Ser Phe Leu Arg Glu His Phe Phe  
 65 70 75 80  
 His Glu Gly Arg Leu Ser Lys Glu Gln Ala Ile Lys Ile Leu Asn Met  
 85 90 95  
 Ser Thr Val Ala Leu Ser Lys Glu Pro Asn Leu Leu Lys Leu Lys Ala  
 100 105 110  
 Pro Ile Thr Ile Cys Gly Asp Ile His Gly Gln Tyr Tyr Asp Leu Leu  
 115 120 125  
 Lys Leu Phe Glu Val Gly Gly Asp Pro Ala Glu Ile Asp Tyr Leu Phe  
 130 135 140  
 Leu Gly Asp Tyr Val Asp Arg Gly Ala Phe Ser Phe Glu Cys Leu Ile  
 145 150 155 160  
 Tyr Leu Tyr Ser Leu Lys Leu Asn Asn Leu Gly Arg Phe Trp Met Leu  
 165 170 175  
 Arg Gly Asn His Glu Cys Lys His Leu Thr Ser Tyr Phe Thr Phe Lys  
 180 185 190  
 Asn Glu Met Leu His Lys Tyr Asp Met Glu Val Tyr Asp Ala Cys Cys  
 195 200 205  
 Arg Ser Phe Asn Val Leu Pro Leu Ala Ala Leu Met Asn Gly Gln Tyr  
 210 215 220  
 Phe Cys Val His Gly Gly Ile Ser Pro Glu Leu Lys Ser Val Glu Asp  
 225 230 235 240  
 Val Asn Lys Ile Asn Arg Phe Arg Glu Ile Pro Ser Arg Gly Leu Met  
 245 250 255

Cys Asp Leu Leu Trp Ala Asp Pro Val Glu Asn Tyr Asp Asp Ala Arg  
                   260                                  265                                  270  
 Asp Gly Ser Glu Phe Asp Gln Ser Glu Asp Glu Phe Val Pro Asn Ser  
                   275                                  280                                  285  
 Leu Arg Gly Cys Ser Phe Ala Phe Thr Phe Lys Ala Ser Cys Lys Phe  
                   290                                  295                                  300  
 Leu Lys Ala Asn Gly Leu Leu Ser Ile Ile Arg Ala His Glu Ala Gln  
                   305                                  310                                  315                                  320  
 Asp Ala Gly Tyr Arg Met Tyr Lys Asn Asn Lys Val Thr Gly Phe Pro  
                                   325                                  330                                  335  
 Ser Leu Ile Thr Met Phe Ser Ala Pro Asn Tyr Leu Asp Thr Tyr His  
                                   340                                  345                                  350  
 Asn Lys Ala Ala Val Leu Lys Tyr Glu Glu Asn Val Met Asn Ile Arg  
                   355                                  360                                  365  
 Gln Phe His Met Ser Pro His Pro Tyr Trp Leu Pro Asp Phe Met Asp  
                   370                                  375                                  380  
 Val Phe Thr Trp Ser Leu Pro Phe Val Gly Glu Lys Val Thr Ser Met  
                   385                                  390                                  395                                  400  
 Leu Val Ser Ile Leu Asn Ile Cys Ser Glu Gln Glu Leu Asp Pro Glu  
                                   405                                  410                                  415  
 Ser Glu Pro Lys Ala Ala Glu Glu Thr Val Lys Ala Arg Ala Asn Ala  
                                   420                                  425                                  430  
 Thr Lys Glu Thr Gly Thr Pro Ser Asp Glu Lys Ala Ser Ser Ala Ile  
                   435                                  440                                  445  
 Leu Glu Asp Glu Thr Arg Arg Lys Ala Leu Arg Asn Lys Ile Leu Ala  
                   450                                  455                                  460  
 Ile Ala Lys Val Ser Arg Met Phe Ser Val Leu Arg Glu Glu Ser Glu  
                   465                                  470                                  475                                  480  
 Lys Val Glu Tyr Leu Lys Thr Met Asn Ala Gly Val Leu Pro Arg Gly  
                                   485                                  490                                  495  
 Ala Leu Ala Arg Gly Thr Glu Gly Leu Asn Glu Thr Leu  
                   500                                  505

## (2) INFORMATION FOR SEQ ID NO:20:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1767 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: unknown

## (ii) MOLECULE TYPE: DNA

## (iii) HYPOTHETICAL: NO

## (iv) ANTI-SENSE: NO

## (vi) ORIGINAL SOURCE:

- (C) INDIVIDUAL ISOLATE: DNA fragment containing  
CNA2deltaC coding sequence

## (ix) FEATURE:

(A) NAME/KEY: CDS  
(B) LOCATION: 262..1767

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ATAGTCTATA ATACGTTTGA TACAGCTAGA TATCGCTAGC GCCAACATTG TCCCCCTCTC	60
TTGATCAATG CTTTTTTTCG GCGGAGACA AATGAGAAAA TGTCCTAAAA ATACCTTTCA	120
TCAAGACTCC TATTTTTCCT TAGAAAAAAC ATATATCCAA CTGGAACAGT ATTAAGCCAA	180
TTGCTACGAT ACAAACAAAA GGAGATATTC CTTCCCTCCC ATAGAGTCAC ACAGGAGCCA	240
GTACTTCTTC TTGAACCCGC A ATG TCT TCA GAC GCT ATA AGA AAT ACT GAG	291
Met Ser Ser Asp Ala Ile Arg Asn Thr Glu	
1 5 10	
CAG ATA AAC GCC GCT ATT AAA ATT ATA GAA AAC AAA ACA GAG CGT CCG	339
Gln Ile Asn Ala Ala Ile Lys Ile Ile Glu Asn Lys Thr Glu Arg Pro	
15 20 25	
CAA TCG TCC ACA ACC CCT ATA GAT TCG AAG GCT AGT ACA GTT GCT GCT	387
Gln Ser Ser Thr Thr Pro Ile Asp Ser Lys Ala Ser Thr Val Ala Ala	
30 35 40	
GCT AAT TCC ACG GCC ACA GAA ACT TCC AGA GAC CTT ACA CAA TAT ACC	435
Ala Asn Ser Thr Ala Thr Glu Thr Ser Arg Asp Leu Thr Gln Tyr Thr	
45 50 55	
CTA GAT GAC GGA AGA GTC GTA TCG ACA AAC CGC AGA ATA ATG AAT AAA	483
Leu Asp Asp Gly Arg Val Val Ser Thr Asn Arg Arg Ile Met Asn Lys	
60 65 70	
GTG CCC GCC ATC ACG TCA CAT GTT CCT ACA GAT GAA GAG CTG TTC CAG	531
Val Pro Ala Ile Thr Ser His Val Pro Thr Asp Glu Glu Leu Phe Gln	
75 80 85 90	
CCC AAT GGG ATA CCT CGT CAC GAA TTC CTA AGA GAT CAT TTC AAG CGC	579
Pro Asn Gly Ile Pro Arg His Glu Phe Leu Arg Asp His Phe Lys Arg	
95 100 105	
GAG GGC AAA TTG TCG GCT GCG CAG GCG GCC AGG ATC GTT ACA CTT GCA	627
Glu Gly Lys Leu Ser Ala Ala Gln Ala Ala Arg Ile Val Thr Leu Ala	
110 115 120	
ACG GAA CTC TTC AGC AAA GAA CCC AAC CTT ATA TCT GTT CCC GCC CCA	675
Thr Glu Leu Phe Ser Lys Glu Pro Asn Leu Ile Ser Val Pro Ala Pro	
125 130 135	
ATC ACA GTT TGC GGT GAT ATC CAT GGC CAG TAC TTT GAC CTT TTG AAG	723
Ile Thr Val Cys Gly Asp Ile His Gly Gln Tyr Phe Asp Leu Leu Lys	
140 145 150	
CTA TTC GAA GTT GGC GGA GAT CCG GCC ACT ACA TCG TAT TTG TTC TTG	771
Leu Phe Glu Val Gly Gly Asp Pro Ala Thr Thr Ser Tyr Leu Phe Leu	
155 160 165 170	
GGA GAC TAT GTC GAC AGA GGG TCC TTT TCG TTT GAG TGT CTT ATT TAT	819
Gly Asp Tyr Val Asp Arg Gly Ser Phe Ser Phe Glu Cys Leu Ile Tyr	
175 180 185	
TTA TAT TCT TTG AAG CTG AAT TTT AAC GAC CAT TTC TGG CTA CTG AGG	867
Leu Tyr Ser Leu Lys Leu Asn Phe Asn Asp His Phe Trp Leu Leu Arg	
190 195 200	

GGT AAC CAC GAA TGT AAG CAT CTA ACG TCA TAT TTC ACT TTC AAA AAT Gly Asn His Glu Cys Lys His Leu Thr Ser Tyr Phe Thr Phe Lys Asn 205 210 215	915
GAA ATG CTG CAC AAG TAC AAT CTA GAT ATT TAC GAG AAA TGC TGC GAA Glu Met Leu His Lys Tyr Asn Leu Asp Ile Tyr Glu Lys Cys Cys Glu 220 225 230	963
TCG TTT AAC AAC TTG CCC CTG GCT GCG TTA ATG AAC GGA CAG TAT CTT Ser Phe Asn Asn Leu Pro Leu Ala Ala Leu Met Asn Gly Gln Tyr Leu 235 240 245 250	1011
TGT GTT CAT GGT GGT ATA TCT CCC GAG TTA AAC TCT TTA CAG GAC ATT Cys Val His Gly Gly Ile Ser Pro Glu Leu Asn Ser Leu Gln Asp Ile 255 260 265	1059
AAC AAC CTA AAT AGA TTC AGG GAG ATT CCC TCT CAT GGC CTG ATG TGT Asn Asn Leu Asn Arg Phe Arg Glu Ile Pro Ser His Gly Leu Met Cys 270 275 280	1107
GAT CTG TTG TGG GCT GAC CCG ATT GAA GAG TAC GAC GAA GTC TTG GAT Asp Leu Leu Trp Ala Asp Pro Ile Glu Glu Tyr Asp Glu Val Leu Asp 285 290 295	1155
AAA GAC TTG ACT GAG GAA GAC ATA GTG AAC TCC AAA ACC ATG GTT CCT Lys Asp Leu Thr Glu Glu Asp Ile Val Asn Ser Lys Thr Met Val Pro 300 305 310	1203
CAT CAT GGC AAG ATG GCA CCT TCA AGG GAT ATG TTT GTC CCA AAC TCA His His Gly Lys Met Ala Pro Ser Arg Asp Met Phe Val Pro Asn Ser 315 320 325 330	1251
GTA AGG GGC TGT TCA TAT GCC TTC ACG TAT CGT GCT GCG TGC CAT TTT Val Arg Gly Cys Ser Tyr Ala Phe Thr Tyr Arg Ala Ala Cys His Phe 335 340 345	1299
CTG CAA GAG ACT GGC CTG TTG TCC ATC ATC AGG GCA CAC GAG GCT CAA Leu Gln Glu Thr Gly Leu Leu Ser Ile Ile Arg Ala His Glu Ala Gln 350 355 360	1347
GAC GCT GGT TAT AGA ATG TAC AAA AAT ACC AAG ACT TTG GGC TTT CCC Asp Ala Gly Tyr Arg Met Tyr Lys Asn Thr Lys Thr Leu Gly Phe Pro 365 370 375	1395
TCT CTT TTG ACC CTT TTC AGT GCG CCT AAC TAC TTG GAC ACC TAC AAT Ser Leu Leu Thr Leu Phe Ser Ala Pro Asn Tyr Leu Asp Thr Tyr Asn 380 385 390	1443
AAT AAG GCT GCC ATA TTG AAA TAC GAA AAT AAT GTT ATG AAT ATC AGA Asn Lys Ala Ala Ile Leu Lys Tyr Glu Asn Asn Val Met Asn Ile Arg 395 400 405 410	1491
CAA TTC AAC ATG ACT CCA CAC CCC TAT TGG TTA CCA GAT TTC ATG GAC Gln Phe Asn Met Thr Pro His Pro Tyr Trp Leu Pro Asp Phe Met Asp 415 420 425	1539
GTT TTC ACG TGG TCC TTG CCA TTT GTT GGT GAA AAA GTT ACA GAG ATG Val Phe Thr Trp Ser Leu Pro Phe Val Gly Glu Lys Val Thr Glu Met 430 435 440	1587
CTT GTC GCA ATT CTA AAC ATC TGT ACT GAA GAT GAG CTG GAA AAC GAC Leu Val Ala Ile Leu Asn Ile Cys Thr Glu Asp Glu Leu Glu Asn Asp 445 450 455	1635
ACC CCC GTC ATT GAA GAA TTA GTT GGT ACC GAT AAA AAA TTG CCA CAA Thr Pro Val Ile Glu Glu Leu Val Gly Thr Asp Lys Lys Leu Pro Gln 460 465 470	1683



71

GCT GGT AAG TCG GAA GCA ACT CCA CAA CCA GCC ACT TCG GCG TCG CCT 1731  
 Ala Gly Lys Ser Glu Ala Thr Pro Gln Pro Ala Thr Ser Ala Ser Pro  
 475 480 485 490

AAA CAT GCT TCC ATT TTA GAT GAC GAA CAT CGA AGG 1767  
 Lys His Ala Ser Ile Leu Asp Asp Glu His Arg Arg  
 495 500

## (2) INFORMATION FOR SEQ ID NO:21:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 502 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Met Ser Ser Asp Ala Ile Arg Asn Thr Glu Gln Ile Asn Ala Ala Ile  
 1 5 10 15  
 Lys Ile Ile Glu Asn Lys Thr Glu Arg Pro Gln Ser Ser Thr Thr Pro  
 20 25 30  
 Ile Asp Ser Lys Ala Ser Thr Val Ala Ala Asn Ser Thr Ala Thr  
 35 40 45  
 Glu Thr Ser Arg Asp Leu Thr Gln Tyr Thr Leu Asp Asp Gly Arg Val  
 50 55 60  
 Val Ser Thr Asn Arg Arg Ile Met Asn Lys Val Pro Ala Ile Thr Ser  
 65 70 75 80  
 His Val Pro Thr Asp Glu Glu Leu Phe Gln Pro Asn Gly Ile Pro Arg  
 85 90 95  
 His Glu Phe Leu Arg Asp His Phe Lys Arg Glu Gly Lys Leu Ser Ala  
 100 105 110  
 Ala Gln Ala Ala Arg Ile Val Thr Leu Ala Thr Glu Leu Phe Ser Lys  
 115 120 125  
 Glu Pro Asn Leu Ile Ser Val Pro Ala Pro Ile Thr Val Cys Gly Asp  
 130 135 140  
 Ile His Gly Gln Tyr Phe Asp Leu Leu Lys Leu Phe Glu Val Gly Gly  
 145 150 155 160  
 Asp Pro Ala Thr Thr Ser Tyr Leu Phe Leu Gly Asp Tyr Val Asp Arg  
 165 170 175  
 Gly Ser Phe Ser Phe Glu Cys Leu Ile Tyr Leu Tyr Ser Leu Lys Leu  
 180 185 190  
 Asn Phe Asn Asp His Phe Trp Leu Leu Arg Gly Asn His Glu Cys Lys  
 195 200 205  
 His Leu Thr Ser Tyr Phe Thr Phe Lys Asn Glu Met Leu His Lys Tyr  
 210 215 220  
 Asn Leu Asp Ile Tyr Glu Lys Cys Cys Glu Ser Phe Asn Asn Leu Pro  
 225 230 235 240  
 Leu Ala Ala Leu Met Asn Gly Gln Tyr Leu Cys Val His Gly Gly Ile  
 245 250 255

Ser Pro Glu Leu Asn Ser Leu Gln Asp Ile Asn Asn Leu Asn Arg Phe  
 260 265 270  
 Arg Glu Ile Pro Ser His Gly Leu Met Cys Asp Leu Leu Trp Ala Asp  
 275 280 285  
 Pro Ile Glu Glu Tyr Asp Glu Val Leu Asp Lys Asp Leu Thr Glu Glu  
 290 295 300  
 Asp Ile Val Asn Ser Lys Thr Met Val Pro His His Gly Lys Met Ala  
 305 310 315 320  
 Pro Ser Arg Asp Met Phe Val Pro Asn Ser Val Arg Gly Cys Ser Tyr  
 325 330 335  
 Ala Phe Thr Tyr Arg Ala Ala Cys His Phe Leu Gln Glu Thr Gly Leu  
 340 345 350  
 Leu Ser Ile Ile Arg Ala His Glu Ala Gln Asp Ala Gly Tyr Arg Met  
 355 360 365  
 Tyr Lys Asn Thr Lys Thr Leu Gly Phe Pro Ser Leu Leu Thr Leu Phe  
 370 375 380  
 Ser Ala Pro Asn Tyr Leu Asp Thr Tyr Asn Asn Lys Ala Ala Ile Leu  
 385 390 395 400  
 Lys Tyr Glu Asn Asn Val Met Asn Ile Arg Gln Phe Asn Met Thr Pro  
 405 410 415  
 His Pro Tyr Trp Leu Pro Asp Phe Met Asp Val Phe Thr Trp Ser Leu  
 420 425 430  
 Pro Phe Val Gly Glu Lys Val Thr Glu Met Leu Val Ala Ile Leu Asn  
 435 440 445  
 Ile Cys Thr Glu Asp Glu Leu Glu Asn Asp Thr Pro Val Ile Glu Glu  
 450 455 460  
 Leu Val Gly Thr Asp Lys Lys Leu Pro Gln Ala Gly Lys Ser Glu Ala  
 465 470 475 480  
 Thr Pro Gln Pro Ala Thr Ser Ala Ser Pro Lys His Ala Ser Ile Leu  
 485 490 495  
 Asp Asp Glu His Arg Arg  
 500

## (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: G4-PCR-A

73

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GCCCTATCGT GCACTCACCG ACGC

24

(2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 24 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: G4-PCR-B

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GTGAAGGCC C TACTGAGCCA GGAG

24

## IT IS CLAIMED:

1. A polypeptide composition comprising a polypeptide effective to enhance immunosuppressive effects of a calcineurin-targeted immunosuppressant by potentiating an interaction of an immunophilin with calcineurin.  
5
2. A composition of claim 1, wherein the polypeptide composition contains a calcineurin interacting (CNI) polypeptide.
- 10 3. A composition of claim 2, wherein the polypeptide composition contains a polypeptide having an amino acid sequence selected from the group consisting of sequences represented by SEQ ID NO:2 and SEQ ID NO:5.
4. A composition of claim 2, wherein the polypeptide has an amino acid sequence of  
15 between 15 and 915 amino acids in length.
5. A composition of claim 2, wherein the polypeptide composition contains a polypeptide having an amino acid sequence comprising the c-terminal 306 amino acids of a CNI protein.  
20
6. An isolated nucleic acid having a sequence encoding a polypeptide of any of claims 1-5.
7. A nucleic acid of claim 6, wherein the nucleic acid has a sequence selected from  
25 the group consisting of nucleic acid sequences represented by SEQ ID NO:3 and SEQ ID NO:6.
8. A method of identifying a small molecule immunosuppressant compound, comprising  
30 constructing a cell-based two hybrid protein-protein interaction assay, wherein one of two fusion hybrid proteins in a cell contains an (A) subunit of calcineurin, and the other of two fusion hybrid proteins contains a CNI polypeptide,  
contacting the cell with a small molecule, and

identifying the small molecule as an immunosuppressant if the molecule potentiates an interaction between the two hybrid proteins.

9. A method of claim 8, wherein the cell is a yeast cell.

5

10. A method of claim 8, wherein one of the two fusion hybrid proteins contains a GAL4 activation domain and the other of two fusion hybrid proteins contains a GAL4 binding domain.

10

11. A method of claim 8, wherein the subunit of calcineurin is selected from the group consisting of yeast CNA1 and yeast CNA2.

12. A method of claim 8, wherein the subunit of calcineurin is an "A" subunit of human calcineurin.

15

13. A method of claim 8, wherein the CNI polypeptide is yeast CNI polypeptide.


14. A method of claim 8, wherein the CNI polypeptide is yeast CN1c polypeptide.

20

15. A method of claim 8, wherein the cell is further modified to cause overexpression of a "B" subunit of calcineurin by said cell.


25

16. A yeast cell carrying a mutation in the naturally-occurring genomic copy of a gene encoding calcineurin-interacting polypeptide, where said mutation prevents expression of a functional calcineurin-interacting polypeptide from said genomic copy.




**Fig. 1B**



**Fig. 2A**



**Fig. 2B**




2/11


**Fig. 3A****Fig. 3B**

SUBSTITUTE SHEET (RULE 26)


3/1.1




**Fig. 4A**



**Fig. 4C**



**Fig. 4B**







**Fig. 4D**

SUBSTITUTE SHEET (RULE 26)



4/11

**Fig. 5A****Fig. 5C****Fig. 5B****Fig. 5D**

SUBSTITUTE SHEET (RULE 26)

5/11




Fig. 6E




Fig. 6C




Fig. 6A




Fig. 6F




Fig. 6D





Fig. 6B


SUBSTITUTE SHEET (RULE 26)

6/11




**Fig. 7A**

**Fig. 7C**




**Fig. 7B**



**Fig. 7D**


**SUBSTITUTE SHEET (RULE 26)**

7/11




**Fig. 7E**

**Fig. 7G**




**Fig. 7F**




**Fig. 7H**

SUBSTITUTE SHEET (RULE 26)


8/11



**Fig. 8A**




**Fig. 8B**




**Fig. 9A**



**Fig. 9C**




**Fig. 9B**




**Fig. 9D**

SUBSTITUTE SHEET (RULE 26)



**Fig. 10**



**Fig. 11**

SUBSTITUTE SHEET (RULE 26)

10/11

tttccagcgg	catatcgtgg	ctttcaccca	acacttcctt	cgagagagtg	cattttacta	61
tgtgaaccaa	tttttcctct	ttttcggttt	gcaagttcac	ctgaaaaact	gcttaacact	121
actagcaatt	gccctattgt	cgtacgagga	ctttgccaaa	tgtattcccc	gctgtttgta	181
gtatatatac	gcagatatat	aatagcgccg	tctttttacc	tctttgagcg	aattgccaaa	241
tattgactct	tttgtcttat	ttcgctatcc	ccatcttata	aaaaatggga	acaactcgtt	301
gaaataagag	acaagcaaca	agaaaagacaa	ccaacagaaa	gttccattcc	gcacaaatac	361
gctggaatcc	catagaatat	tgcttggtcc	tctatgacta	catgctccaa	ttcaatacac	421
aaaatgatac	tgtagctcca	gtgtttccca	tggagcaaga	tataaatgca	gcacctgatg	481
ccgtcccact	ggtgcagaca	acaacactac	aagtctttgt	aaagcttgcc	gaacccatag	541
tgtttttaaa	aggatttgaa	actaacggac	tgtctgaaat	agcccccagt	atcttacgag	601
gatctcttat	cgtcaggggtg	ttgaaaccga	ataaattaaa	aagtatatcg	ataaccttca	661
aaggaatatc	cagaacagag	tggccgggaag	gtataccacc	gaagagagaa	gaatttttcag	721
atgttgaaac	tggtgtcaat	cacacatggc	cattttatca	ggcgggatgac	ggcatgaatt	781
ctttcacctt	agaacatcac	agctcaaata	attcgtccaa	tcgccccatc	atgagcgatg	841
aagattatct	tactgaaaaa	agcgggtgctt	cagtatatat	cccaccaacc	gctgaacccc	901
ctaaagataa	tagcaatcta	agtctggatg	cctatgagcg	caactcattg	tcattccgata	961
atttgagtaa	caagccagta	tcaagtgatg	tttcccatga	cgacagtaaa	ctgttggcta	1021
ttcaaaagac	accattacca	tcattctagtc	gaagaggatc	ggtaccggca	aattttcacg	1081
gtaactcttt	gtcacctcat	accttcatat	ctgatttggt	cacaaaaaca	ttcagtaata	1141
gtggcgctac	tccaagtcct	gagcaagagg	ataactatct	tacaccatcc	aaagattcta	1201
aagaagtttt	tatttttcga	ccggggcgatt	atattttacac	ttttgaacag	ccaatatcgc	1261
aatcttatcc	agaaagtata	aaagccaatt	ttggttccgt	ggagtataaa	ctgtcaatag	1321
acatagagag	gttttgcgca	ttcaaatcaa	ctatacatat	tcaattaccc	atcaaatgctg	1381
taaggcttcc	ttctgatgga	tccgtagaag	agactgaagc	tattgcaatt	tccaaggact	1441
ggaaagatct	tcttcattat	gacgtggtaa	ttttctcgaa	agagatcggt	ttgaatgcat	1501
ttttacccat	cgattttccat	ttcgctcctc	tagataaagt	tactctgcat	cgtattagaa	1561
tttatctaac	agagtctatg	gaatacactt	gtaatagtaa	tggaaatcac	gagaaggctc	1621
gtagattaga	gccaactaaa	aagtttctgt	tggctgaaca	taacggctct	aaactgcctc	1681
atataccagc	tggttcgaat	cctttgaagg	ctaaaaatag	agggaaacatc	ctcttggatg	1741
aaaaatccgg	cgatctagtt	aacaaagatt	ttcagttcga	ggtgtttgtc	ccaagcaagt	1801
ttacaaacag	tatacgggta	caccctgata	caaattatga	taaaatcaaa	gcccaccatt	1861
ggataaaaaat	ttgccttcgt	ctttccaaga	agtacgggga	caatagaaaa	cattttcgaaa	1921
taagtattga	ttctccaatc	catattttaa	atcaactatg	ctcacacgcg	aatactttgc	1981
taccgagcta	cgagagtcac	ttccagtatt	gtgatgaaga	tggttaatttc	gcaccagcat	2041
cagatcaaca	aaattacgca	agtcatcatg	attccaatat	tttcttccca	aaagaagttc	2101
tttcgtctcc	cgttcttttca	cctaacgtgc	agaagatgaa	cattagaata	ccgtctgcat	2161
ttccagtagt	gcgtaataga	gctgaaagcg	taaagaaaag	caagtcagat	aatacctcca	2221
agaagaatga	tcaaagtagc	aatgtcttcg	catccaaaca	gctggctcgca	aacattttata	2281
agcccaatca	gattccaaga	gaattaaact	ctcctcaggc	gttaccatta	tcgccccatca	2341
cctcaccaat	tctcaattac	caaccattat	caaactcccc	gcctccagat	tttgattttg	2401
atctagctaa	gcgcggcgca	gccgattctc	atgctattcc	tgtggatcct	ccatcatatt	2461
ttgatgtatt	aaaggccgat	gggattgaat	tgccatacta	cgatacaagt	tcattctaaaa	2521
ttcctgaact	aaaactaaac	aatcttagag	agacattggc	cagcattgag	gaggactcat	2581
tcaatgggtg	gtctcaaatt	gatgacttat	ccgacgaaga	tgacaatgat	ggcgatatag	2641
catctgggtt	caacttcaag	ctgtcaacca	gtgctccgag	tgagaacggt	aattcacaca	2701
ctcctatttt	gcagtcttta	aacatgagtc	ttgatgggag	aaaaaaaaat	cgtgccagtc	2761
tacacgcaac	atcagtgtta	cctagtacaa	taagacagaa	caatcagcat	ttcaatgaca	2821
taaaccagat	gctagggcag	agtgacgaag	atgcctttcc	caaaagccaa	tcattaaatt	2881
tcaataagaa	actaccaata	cttaaaatta	atgataacgt	catacaatca	aacagcaata	2941
gtaataacag	agttgataat	ccagaagata	cagtggaattc	ttcagtcgat	attacagcat	3001
tttatgctcc	aagaattgtca	tcagattcca	aatttgattg	ggaggtaagc	aagaacccatg	3061
ttgaccagc	agcctactcg	gttaacggtg	tacattgaaa	ccgtgtactg	gacgacttta	3121
agaaagcatt	tcgcgaaaag	agaaaataag	tacatttttt	tcatttctcg	acagaattgc	3181
taccatttta	ctttgtgtcc	tgtgattcaa	tagtgtaaaa	tatattggac	attttatag	3241
atacaaatat	acaccatcaa	tctatacatc	catatcactt	gtcgtaaaaga	tatccctttt	3301
taatagtaca	gcgattaaaa	aaataacatg	attaacgt			

Fig. 12

11/11

MLQFNTENDTVAPVFPMEQDINAAPDAVPLVQTTTLQVFVKLAEPVFLKGFETNGLSEIAPSILRGSLIVR  
VLKPNKLKSSITFKGISRTEWPEGIPPKREEFSDVETVVNHTWPFYQADDGMNSFTLEHHSSNNSSNRPSM  
SDEDYLLEKSGASVYIPPTAEPKDNSNLSDAYERNLSSDNLNKPVSSDVSHDDSKLLAIQKTPLPSSS  
RRGSVPANFHGNSLSPHTFISDLFTKTFSNSGATPSPEQEDNYLTPSKDSKEVFIFRPGDYIYTFEQPISQS  
YPESIKANFGSVEYKLSIDIERFGAFKSTIHTQLPIKVVRLPSDGSVEETEAIAISKDWKDLLHYDVVIFSK  
EIVLNAFLPIDFHFAPLDKVTLHRIRIYLTESMEYTCNSNGNHEKARRLEPTKKFLLAEHNGPKLPHIPAGS  
NPLKAKNRGNILLDEKSGDLVNKDFQFEVFVPSKFTNSIRLHPDTNYDKIKAHHWIKICLRLSKKYGDNRRKH  
FEISIDSPIHILNQLCSHANTLLPSYESHFQYCEDGNGFAPAADQQNYASHHDSNIFFPKEVLSSPVLSPNV  
QKMNI<sup>→</sup>RIPSDLPVVRNRAESVKKSKSDNTSKKNDQSSNVFASKQLVANIYKPNQIPRELTS<sup>→</sup>QALPLSPITS  
PILNYQPLSNSPPPDFDFDLAKRGAADSHAIPVDPPSYFDVLKADGIELPYYDTSSSKIPELKLNKSRETLA  
SIEEDSFNGWSQIDDLSDEDDNDGDIASGFNFKLSTSAPSENVNSHTPILQSLNMSLDGRKKNRASLHATSV  
LPSTIRQNNQHFNDINQMLGSSDEDAFPKSQSLNFKKLPIKINDNVIQSNSNSNNRVDPEDTVDSVSDI  
TAFYDPRMSSDSKFDWEVSKNHVDPAAYSVNVASENRVLDDFKKAFREKRK

**Fig. 13**



## INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US 95/13580

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C12N15/31 C07K14/395 G01N33/68 C12N15/62 C12Q1/00  
C12N1/16 //(C12N1/16,C12R1:865)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C12N C07K G01N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Swissprot Database entry YK01_YEAST Accession number P36117; June 01, 1994 Duesterhoeft A. et al.: 'Hypothetical 102.5 kD protein in YPT52-GCN3 intergenic region.'	1-5
X	Emfun Database entry Scykr021w Accession number Z28246; May 10, 1994 Duesterhoeft A. et al.: 'S. cerevisiae chromosome XI reading frame ORF YKR021w.'	6,7
P,X	MOLECULAR BIOLOGY OF THE CELL, 5 (SUPPL.). 1994. 141A., HUANG L ET AL 'A novel protein that interacts with calcineurin in vivo' see abstract 818	1,2,6, 8-15



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*A\* document member of the same patent family

Date of the actual completion of the international search

21 March 1996

Date of mailing of the international search report

02.04.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Espen, J

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 95/13580

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>CURR BIOL, 2 (1). 1992. 18-20., CYERT M S 'IMMUNOSUPPRESSANTS HIT THE TARGET' see the whole document -----</p>	